Short prokaryotic Argonautes provide defence against incoming mobile genetic elements through NAD + depletion

Nature Microbiology · 2021 · 10.1101/2021.12.14.472599

8
Short prokaryotic Argonautes provide defence against incoming mobile genetic
1
elements through NAD+ depletion
2
Mindaugas Zaremba1*, Donata Dakineviciene1, Edvardas Golovinas1, Evelina Zagorskait1, Edvinas
3
Stankunas1, Anna Lopatina2, Rotem Sorek2, Elena Manakova1, Audrone Ruksenaite1, Arunas
4
Silanskas1, Simonas Asmontas1, Algirdas Grybauskas1, Ugne Tylenyte1, Edvinas Jurgelaitis1, Rokas
5
Grigaitis1#, Kstutis Timinskas1, eslovas Venclovas1, Virginijus Siksnys1*
6
7
1Institute of Biotechnology, Life Sciences Center, Vilnius University, Saultekio av. 7, LT-10257,
8
Vilnius, Lithuania.
9
2Department of Molecular Genetics, Weizmann Institute of Science, Rehovot 7610001, Israel.
10
*To whom correspondence should be addressed. Tel: +370-5-2234357; Fax: +370-5-2234367;
11
Email: zare@ibt.lt. Correspondence may also be addressed to Virginijus Siksnys. Tel: +370-5-
12
2234359; Fax: +370-5-2234367; Email: siksnys@ibt.lt.
13
The first 4 authors contributed equally to this manuscript.
14
Present address: Edvinas Stankunas, Max F. Perutz Laboratories, Medical University of Vienna,
15
Vienna Biocenter (VBC), Dr. Bohr-Gasse 9/3, 1030 Vienna, Austria.
16
Anna Lopatina, Division of Microbial Ecology, Department of Microbiology and Ecosystem
17
Science, Center for Microbiology and Environmental Systems Science, University of Vienna,
18
Vienna, Austria.
19
Rokas Grigaitis, VU LSC-EMBL Partnership for Genome Editing Technologies, Life Sciences
20
Center, Vilnius University, Saultekio av. 7, LT-10257, Vilnius, Lithuania.
21
22
23

.
CC-BY-NC-ND 4.0 International license
made available under a
(which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is The copyright holder for this preprint
this version posted August 25, 2022.
;
https://doi.org/10.1101/2021.12.14.472599
doi:
bioRxiv preprint
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
2
Argonaute (Ago) proteins are found in all three domains of life. The so-called long Agos are composed of four major domains (N, PAZ, MID, and PIWI) and contribute to RNA silencing in eukaryotes (eAgos) or defence against invading mobile genetic elements in prokaryotes (pAgos). The majority (~60%) of pAgos identified bioinformatically are shorter (comprised of only MID and PIWI domains) and are typically associated with Sir2, Mrr or TIR domain-containing proteins. The cellular function and mechanism of short pAgos remain enigmatic. Here, we show that Geobacter sulfurreducens short pAgo and the NAD +-bound Sir2-protein form a stable heterodimeric complex. The GsSir2/Ago complex presumably recognizes invading plasmid or phage DNA and activates the Sir2 subunit, which triggers endogenous NAD + depletion and cell death, and prevents the propagation of invading DNA. We reconstituted NAD + depletion activity in vitro and showed that activated GsSir2/Ago complex functions as a NADase that hydrolyses NAD + to ADPR. Thus, short Sir2-associated pAgos provide defence against phages and plasmids and underscores the diversity of mechanisms of prokaryotic Agos.
38
Being at the core of RNA interference eAgos are involved in the regulation of gene expression, silencing of mobile genome elements, and defence against viruses 1,2. The best-studied hAgo2 uses small RNA molecules as guides for target RNA recognition, and eAgos are similar both structurally and mechanistically 2–5. Monomeric eAgos are composed of four major N, PAZ (Piwi/Argonaute/Zwille), MID (Middle), and PIWI (P-element Induced Wimpy testis) domains (Fig. 1A) and share a bilobed structure, where the N- and C-terminal lobes are formed by conserved N/PAZ and MID/PIWI domains, respectively 4–7. The N-domain acts as a wedge that separates guide and target strands 3,7, while the MID and PAZ domains bind, respectively, the 5'- and 3'-terminus of the guide RNA, located between the N- and C-terminal lobes 4,6. eAgos can slice target RNA through endonucleolytic cleavage by the PIWI domain or inhibit translation through RNA binding by the catalytically inactive eAgos that may also trigger RNA decay by auxiliary cellular nucleases 2,4,5.

pAgos are quite widespread and are present in 9% of sequenced bacterial and 32% of archaeal genomes^{6. To date, more than ~1000 pAgos have been identified bioinformatically, revealing a striking diversity. Moreover, pAgos are often associated with additional putative nucleases, helicases, and DNA binding proteins that are not linked to eAgos^{8. pAgos are divided into full-length or long pAgos (~40%) sharing conserved N, PAZ, MID, and PIWI domain architecture with eAgos, and short pAgos (~60%) composed only of MID and PIWI domains (Fig. 1A)8,9. Long pAgos are relatively well-characterized both structurally and functionally10,11 and, similarly to eAgos, contain either catalytically active or inactive PIWI domain. Some long pAgos with the catalytically active PIWI, as exemplified by CbAgo and TtAgo, use DNA guides to target and cleave DNA providing defence against invading phages or plasmids, or contributing to chromosome segregation after replication, respectively12–14. Meanwhile, long RsAgo with an inactive PIWI domain

guided by small RNA is thought to mobilize an unknown cellular nuclease(s) for degradation of invading plasmids and mobile genetic elements9,15. Interestingly, long KmAgo can use both DNA and RNA guides to target DNA and RNA in vitro, albeit with different efficiencies16,17. In contrast to long pAgos, all short pAgos possess a catalytically inactive PIWI domain and are typically associated with proteins containing a domain initially thought to be analogous to PAZ (APAZ)18. Subsequently, however, it was proposed that APAZ may actually be homologous to the N-terminal domains (N and L1) of Ago11,19. APAZ-containing proteins are often fused to Sir2 (Silent informator regulator 2), Mrr nucleases or TIR (Toll-Interleukin-1 Receptor) domains8,18. About half of all identified short pAgos are associated or fused into a single-chain

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

protein with Sir2-APAZ proteins 8. The Sir2 domain-containing proteins are widely distributed in all domains of life and perform protein deacetylation or ADP-ribosylation functions using NAD + as a co-factor 20,21. Particularly, bacterial Sir2 proteins are involved in many cellular processes including transcription, translation, carbon and nitrogen metabolism, virulence and resistance to stress 21. Despite the fact that short pAgos, half of which are associated with Sir2 proteins, make up the majority of all pAgos, their function in the cell and in vitro remains to be established.

77 Results

In this study, we aimed to explore whether short pAgos can act as prokaryotic defence systems against viruses or plasmids. To this end, we selected two short pAgos, GsSir2/Ago from Geobacter sulfurreducens and CcSir2/Ago from Caballeronia cordobensis, each encoded in a putative operon together with a Sir2 domain protein, and PgSir2-Ago from Paraburkholderia graminis, representing a fusion of Sir2 and pAgo (Fig. 1A). The coding regions of the Sir2 and pAgo proteins in GsSir2/Ago and CcSir2/Ago systems overlap by 11 and 8 bp, respectively, indicating that they belong to the same operon (Supplementary Text). Next, we engineered heterogeneous E. coli cells by cloning GsSir2/Ago and CcSir2/Ago genes, or a single gene, encoding PgSir2-Ago into pBAD expression vectors under a PBAD promoter (Supplementary Table 1) and challenged them with phages or plasmids.

Sir2/Ago systems provide defence against phages. To test whether the GsSir2/Ago system provides defence against phages, we challenged E. coli host carrying the GsSir2/Ago system with a set of six E. coli phages spanning four morphological families including Podoviridae (T7), Siphoviridae (lambda-vir, SECphi27, SECphi18), Myoviridae (T4) and Microviridae (SECphi17, ssDNA phage) (Supplementary Table 1). We measured the efficiency of plating (EOP) of these phages with and without L-arabinose induction of the GsSir2/Ago system. The system showed protection against two out of six phages – lambda-vir (~100-fold) and SECphi27 (~1000-fold) (Fig. 1C).

To probe the role of individual Sir2 and Ago proteins in antiviral defence, we engineered E. coli cells, carrying mutant variants of either the Sir2 or the Ago protein of the GsSir2/Ago system and performed small drop plaque assays using the lambda-vir and SECphi27 phages. In Sir2 variants, the highly-conserved D230 residue, presumably involved in NAD binding was replaced by alanine (Extended Data Fig. 1C)18. Phage challenge assay revealed that the D230A mutation completely abolished defence against both phages (Fig. 1D). The Ago protein is catalytically inactive due to the active site mutations in the PIWI domain, therefore, to obtain a binding-deficient Ago variant, we fused a bulky 29 aa His-StrepII-His-tag (HSH-tag) at the C-terminus that is important for nucleic acid binding in other Ago 3,6,8,22. We found that Ago C-terminal modification abolished protection from both phages (Fig. 1D). Thus, both the Sir2 and the Ago proteins are required for protection against phages by the GsSir2/Ago system.

Additionally, we probed the ability of homologous CcSir2/Ago and PgSir2-Ago systems to restrict lambda-vir and SECphi27 phages (Fig. 1E). After L-arabinose induction, the PgSir2-Ago system showed ~500-fold protection against lambda phage and ~400-fold against SECphi27, while CcSir2/Ago system showed ~1000-fold

protection against lambda and no protection against SECphi27 phage. Thus, homologous Sir2/Ago systems from three phylogenetically distant bacteria showed protection against phage infection, albeit with different efficiency and specificity.

Next, E. coli MG1655 cells either lacking or containing the GsSir2/Ago system were infected in liquid culture with lambda phage at a multiplicity of infection (MOI) 0.05, 0.5 and 5 (Fig. 1F). At

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

high MOI where, on average, a single bacterial cell is infected by a single phage, the culture collapses, while at low MOI the culture survives. This phenotype implies that GsSir2/Ago-mediated defence triggers cell death approximately at the same time when the phage-induced lysis occurs.

Sir2/Ago systems interfere with plasmid transformation. Further, to test whether heterologous expression of the GsSir2/Ago, CcSir2/Ago and PgSir2-Ago systems in E. coli cells (strains BL21-AI and DH10B) provides a barrier for plasmid transformation, four plasmids (pCDF, pCOLA, pACYC184, and pRSF) with different ori regions and copy numbers were used in a plasmid interference assay (Supplementary Table 1, Fig. 2, and Extended Data Fig. 2). We found that GsSir2/Ago system prevented only pCDF plasmid transformation reducing its efficiency nearly ~100-fold (Fig. 2B). Next, we tested pCDF transformation efficiency in E. coli cells expressing the GsSir2/Ago mutants (Fig. 2C). Both Sir2 D230A mutation and pAgo HSH-tag modification that impaired phage restriction also abolished plasmid interference making cells permissive for pCDF plasmid transformation (Fig. 2C). CcSir2/Ago system like GsSir2/Ago provided resistance only for pCDF plasmid transformation (Extended Data Fig. 2B) while cells carrying the single-chain PgSir2-Ago system was permissive for transformation of all four plasmids (Extended Data Fig. 2B).

Interestingly, although the GsSir2/Ago system in E. coli interfered with the pCDF plasmid transformation, pCOLA plasmid that differs mainly in the ori region and antibiotic resistance gene was permissive. To test whether the ori region determines differences in the transformation efficiency between pCDF and pCOLA plasmids, we swapped the ori sequences of pCDF and pCOLA (Fig. 2D). The pCDF plasmid with ColA ori instead of CloDF13 became permissive in E. coli cells expressing the GsSir2/Ago system, whereas transformation of pCOLA plasmid bearing CloDF13 ori instead of ColA ori was prevented. These results indicate that CloDF13 ori is a key element that controls plasmid transformation efficiency in E. coli cells expressing GsSir2/Ago system.

The plasmid interference by the GsSir2/Ago system could be due to either the plasmid entry exclusion, replication inhibition or plasmid degradation. To eliminate the possible role of the plasmid entry barriers on the pCDF plasmid transformation efficiency, we engineered heterogeneous E. coli cells carrying two plasmids: pBAD plasmid providing carbenicillin (Cb) resistance and expressing GsSir2/Ago (or its mutants) under control of PBAD inducible promoter and the pCDF plasmid providing streptomycin (Str) resistance, and tested cell viability in the presence or absence of the inducer. In this case, the pCDF plasmid is already in the cell and provides streptomycin resistance, however antibiotic resistance should be lost if the plasmid is restricted after GsSir2/Ago expression. In the absence of induction, cell viability of E. coli cells carrying wt GsSir2/Ago (or its mutants) and an empty pBAD vector (Extended Data Fig. 2E) was identical. In the presence of the inducer the viability of cells expressing the wt GsSir2/Ago system, but not its mutants, significantly decreased (Fig. 2E), indicating that GsSir2/Ago interferes with the pCDF plasmid already present in the cell. Notably, a decrease in cell viability is observed in E. coli BL21-AI cells (Tc-resistant) without the cell selection for Str and Cb resistance. It cannot be excluded that upon recognition of pCDF the GsSir2/Ago system becomes activated and triggers cell death. A similar cell death phenotype triggered by the GsSir2/Ago has been observed during phage infection in liquid cultures (Fig. 1F). Taken together, these data show that the GsSir2/Ago system acts as a defence system against phages and plasmids via cell death or suicidal mechanism.

Short pAgo and Sir2 form a stable heterodimeric complex. To characterize the Sir2/Ago systems biochemically, we aimed to express individual Sir2 and Ago proteins in E. coli. The GsSir2 and CcSir2 proteins (but not PgSir2-Ago) were expressed and purified by chromatography (Fig. 3A and Extended Data Fig. 3). The N-terminal His-tagged GsSir2 and CcSir2 proteins co-expressed with Ago proteins co-purified on the Ni +-affinity column (Extended Data Fig. 3), indicating that
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Sir2 and pAgo proteins form a stable complex. We failed to express Sir2 and Ago proteins individually, suggesting that they form an obligatory Sir2/Ago complex. Sir2/Ago complex. Functionally compromised GsSir2(D230A)/Ago and GsSir2/Ago-HSH variants also formed a complex indicating that while the introduced mutations abolished the activity in vivo, they did not affect the protein complex structure (Fig. 2C, Extended Data Fig. 3C). Further analysis of the oligomeric state of GsSir2/Ago and CcSir2/Ago complexes in solution using SEC-(MALS), mass photometry and small-angle X-ray scattering (SAXS) showed that heterodimeric complexes are formed in the wide range of protein concentrations (from 20 nM to 6.5 M) (Fig. 3B-D, and Extended Data Fig. 4). According to the SAXS data, the heterodimeric GsSir2/Ago complex acquires a notably asymmetric shape (Fig. 3D, Extended Data Fig. 4, Supplementary Table 3) that is consistent with a structural model of the heterodimer (Supplementary Text). In summary, the results show that pAgos and associated Sir2 proteins encoded by a single operon form a stable heterodimeric complex.
Sir2/Ago prefers an RNA guide to bind a DNA target. Long pAgos use ssDNA and/or ssRNA guides to recognize their complementary DNA and/or RNA targets 10. To establish guide preference of Gs and Cc Sir2/Ago, we analysed by EMSA binding of single- or double-stranded DNA or RNA (Extended Data Fig. 6). Both Sir2/Ago heterodimers showed a strong preference for ssDNA and ssRNA binding. RNA/DNA heteroduplex was bound with an intermediate affinity, while dsRNA or dsDNA showed only weak binding (Extended Data Fig. 6A-B,E, Table 1). Interestingly, neither the 5'-terminal phosphate nor the 3'-OH end or Mg 2+ ions were required for ssDNA binding as both Gs and Cc complexes bound the circular ssDNA and linear oligonucleotides with similar affinity (Extended Data Fig. 6A-B). As expected, Gs and Cc Sir2/Ago containing the inactivated PIWI domains showed no cleavage activity for any NA substrate tested (Extended Data Fig. 6F). In summary, EMSA experiments suggest that in vitro ssRNA or ssDNA are preferable GsSir2/Ago guides.
Next, we analysed ssDNA or ssRNA target binding by the binary GsSir2/Ago complexes preloaded with either ssRNA or ssDNA guides. In a separate set of experiments, reaction mixture also contained heparin, a competitor of nucleic acid binding (Fig. 4A, Table 1). The binary GsSir2/AgossRNA complex showed ~10-fold better binding to the ssDNA than ssRNA, and heparin addition had only a little effect (~4-fold decrease) on binding affinity in this case. The GsSir2/Ago-ssDNA binary complex bound to the matching ssDNA target with affinity similar to the GsSir2/AgossRNA binary complex, however, heparin addition abolished binding (Extended Data Fig. 6A and D). Furthermore, the binary GsSir2/Ago-ssRNA complex bound to the complementary ssDNA target ~200-fold better than apo-GsSir2/Ago bound pre-annealed gRNA/tDNA heteroduplex indicating that GsSir2/Ago requires binding of the RNA guide first to interact with the DNA target (Fig. 4A, Table 1). The binding affinity of the functionally compromised in vivo GsSir2(D230A)/Ago mutant was similar to the wt, while the binding affinity of the GsSir2/Ago-HSH mutant was slightly (~7-fold) weaker (Fig. 4A, Extended Data Fig. 6C, Table 1). Charge reversal mutations of positively charged residues that are involved in the interactions with the guide and target NAs according to the GsSir2/Ago model abolished both the target NA binding and the pCDF plasmid interference (Extended Data Fig. 2F and G, Extended Data Fig. 3B and Extended Data Fig. 6C). Taken together, our data suggest that GsSir2/Ago uses ssRNA as a guide for the recognition of a ssDNA target.
To identify NAs, bound by GsSir2/Ago in vivo, we purified the GsSir2/Ago-NA complex from E. coli transformed with the pBAD_GsSir2/Ago expression vector and the pCDF target plasmid, extracted NAs and

subjected them to sequencing. Subsequent analysis revealed that GsSir2/Ago is associated with small (predominantly 21 nt) RNAs with or without the 5'-phosphate (Fig. 4B and C). Differently from other Argonaute proteins that show base selectivity for the first nucleotide at
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

the 5'-end of the guide 13-15,23, GsSir2/Ago-associated small RNAs show preference for the 5'-AU dinucleotide (Fig. 4D). This preference is more pronounced for small RNAs containing 5'-phosphate, implying that GsSir2/Ago uses as a guide small RNAs containing the 5'-AU dinucleotide. Most co-purified small RNAs (~95%) matched the E. coli genome, whereas the smaller fraction (~5%) originated from the pBAD_GsSir2/Ago and pCDF plasmids (Supplementary File 1). Interestingly, small plasmid-borne RNAs that matched CloDF13 and ColE1 ori regions of corresponding plasmids, were noticeably enriched (Fig. 4E). Taken together, RNA-seq data suggest that GsSir2/Ago could use small 5'-AU-RNAs originating from the invader transcripts (e.g., pCDF ori region) as guides to target the invaders' DNA.

The GsSir2/Ago complex binds NAD + and causes its depletion. Computational analysis of Sir2 domains showed that they possess a conserved NAD +-binding pocket (Extended Data Fig. 1C and Extended Data Fig. 5). To determine whether Sir2 domains can indeed bind endogenous NAD +, purified GsSir2/Ago and CcSir2/Ago complexes were heat-treated, protein aggregates removed by centrifugation, and the supernatant analysed by MS-HPLC (Fig. 5 and Extended Data Fig. 7). The quantitative analysis showed that both the GsSir2/Ago and CcSir2/Ago complexes co-purified with bound endogenous NAD + in approx. 1:1 molar (Sir2:NAD +) ratio (Fig. 5A, Extended Data Fig. 3, Extended Data Fig. 7). However, in the case of the functionally inactive GsSir2(D230A)/Ago mutant, only 0.6% of all complexes were NAD +-bound indicating that the mutation severely compromised NAD + binding by the Sir2 domain (Fig. 5A). NAD + binding by the GsSir2 subunit and the similarity of the GsSir2 to the N-terminal NADase of the ThsA protein from the anti-phage Thoiris system 24 prompted us to investigate the level of endogenous NAD + in the presence of the induced GsSir2/Ago system and its target pCDF plasmid. In these experiments the corresponding E. coli cells were lysed, proteins were removed, and the amount of NAD + in the supernatant was examined by MS-HPLC. When the wt GsSir2/Ago expression was induced in the presence of pCDF plasmid, NAD + was depleted (~120-fold decrease), whereas in the case of the functionally inactive GsSir2(D230A)/Ago and GsSir2/Ago-HSH mutants the level of endogenous NAD + was similar to that of the empty pBAD vector (Fig. 5B, Extended Data Fig. 7). It should be noted that a significant ~30-fold decrease of the NAD + level was observed when the wt GsSir2/Ago expression was induced even in the absence of pCDF suggesting that the heterologously expressed system may be toxic to the cells resulting in their slower growth (Fig. 5B, Extended Data Fig. 7). To test the hypothesis that the activated GsSir2/Ago system similarly to the Thoiris anti-phage system depletes the endogenous NAD + through hydrolysis or cyclization, we attempted to identify possible products (ADPR, cADPR, AMP, cAMP, ADP, cADP, nicotinamide, and adenine) using MS-HPLC albeit without success. It is possible that NAD + conversion products were not detected since in the cell they were processed to other reaction intermediates or due to ion suppression during MS analysis of the cell lysates.

Next, we investigated whether the NAD + depletion activity detected in cells could be reconstituted in vitro. To this aim we mixed the binary GsSir2/Ago-gRNA complex with ssDNA and monitored NAD + concentration using a commercial kit (Fig. 5C). We found that NAD + concentration decreased when wt GsSir2/Ago-gRNA complex was added to the complementary ssDNA target, however, no changes were observed in the case of non-complementary ssDNA. Mutations in the Sir2 domain (D230A) or the APAZ/Ago part (GsSir2/Ago-HSH, GsSir2APAZ/Ago and GsSir2/AgoPIWI mutants) compromised NAD + depletion (Fig. 5C). MS analysis revealed that GsSir2/Ago-gRNA complex in the presence of the complementary ssDNA hydrolyses NAD + to ADPR (Fig. 5D,E) similarly to the Thoeris anti-phage system 24. Taken together, these results demonstrate that GsSir2/Ago functions as a NADase that becomes activated upon target DNA binding.

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

258 Discussion
Association of Sir2-like domains with short pAgo proteins has been identified bioinformatically in the pioneering Makarova et al. paper 18. It has been speculated that Sir2-domain proteins can act as nucleases, however, the structure and function of Sir2 proteins so far have not been elucidated. Here, we show that the APAZ-containing Sir2 and short pAgo proteins form a heterodimeric complex (Fig. 3) similarly as in the case of a short pAgo and a Mrr nuclease domain-containing protein 25. Furthermore, our structure modelling results show that the APAZ region of Sir2 proteins shares similarity with the N, L1 and L2 domains of canonical Agos substantiating previous sequence-based predictions 11,19 At the same time, Sir2 proteins entirely lack the PAZ domain (Fig. 1A, Extended Data Fig. 5A). Thus, apparently both split and single-chain Sir2/Ago systems evolved from long pAgos by the loss of the PAZ domain and acquisition of the Sir2 domain.
Next, we provide the experimental evidence that the Sir2/Ago complex functions as a defence system against invading phages and plasmids (Fig. 1 and Fig. 2). Intriguingly, plasmid interference assay using four plasmids with different replicons (Extended Data Fig. 2) revealed that GsSir2/Ago system prevents transformation only of the pCDF plasmid that contains CloDF13 ori (Fig. 2), suggesting that GsSir2/Ago may recognize specific replicon elements or structures. Indeed, ori swap between pCDF and permissive pCOLA plasmid made the latter sensitive to GsSir2/Ago interference. We show here that GsSir2/Ago co-purifies from E. coli cells together with small (predominantly 21 nt long) RNAs that preferentially contain the 5'-AU dinucleotide (Fig. 4B-D). Interestingly, a fraction of small RNAs that originates from pCDF CloDF13 and pBAD ColE1 or regions is enriched (Fig. 4E). ColE1-like origins, including CloDF13, use two small RNAs (RNAI and RNAII) for priming of the replication that involves the R-loop intermediate26-28. It is tempting to speculate that GsSir2/Ago is able to bind nucleic acids of different length (Fig. 4B,C) and preferentially binds ori-associated small RNAs that can be subjected to further processing by cellular RNases to produce ~21 nt gRNAs similarly to long RsAgo that shares an inactivated PIW domain with GsSir2/Ago15.
We further show that in vitro the reconstituted wt GsSir2/Ago-gRNA complex becomes activated after binding the complementary DNA target and triggers NAD + hydrolysis generating ADR (Fig. 5C-E). It is likely, that in the E. coli cells the APAZ/Ago part in the GsSir2/Ago complex guided by the ori-associated RNA guides could bind to the complementary plasmid or phage DNA target activating the Sir2 effector domain that depletes endogenous NAD + leading to the cell death, thereby restricting plasmid and phage propagation (Fig. 5F, Supplementary text, the accompanying paper by Garb et al., 202129). A similar anti-phage defence mechanism based on NAD + exhaustion has been shown for the Thoeris and the Pycsar systems, CBASS (Cyclic Oligonucleotide-Based Signaling System) and DSR (Defense-Associated Sirtuins)29-31. In the Thoeris system of Bacillus cereus MSX-D12, the Sir2 domain is similar to that of the GsSir2/Ago system and performs the hydrolysis of NAD + to ADR and nicotinamide31 like the GsSir2/Ago system. Further structural and biochemical studies are underway to establish the structure of the heterodimeric Sir2/Ago complex and the mechanism of Sir2 domain activation that triggers NAD + hydrolysis.
297 Methods
Oligonucleotides used in this work. All synthetic DNA oligonucleotides used for cloning and site-specific mutagenesis were purchased from Metabion (Germany) and are listed in Supplementary Table 2.
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license

Cloning and mutagenesis. A whole operon of the GsSir2/Ago system, composed of the Sir2 (GSU1360, NP_952413.1) and Ago (GSU1361, NP_952414.1) encoding genes, was amplified by PCR from the genomic DNA of Geobacter sulfurreducens Caccavo (ATCC No. 51573, LGC Standards cat#51573D-5) using the oligonucleotides MZ-239 and MZ-240 (Supplementary Table 2), respectively. The resulting DNA fragment was digested by Eco31I (ThermoFisher cat#FD0293) and XhoI (ThermoFisher cat#FD0694) and cloned into pBAD/HisA expression vector (ThermoFisher cat#V43001) pre-cleaved with NheI (ThermoFisher cat#FD0973) and XhoI and dephosphorylated using FastAP (ThermoFisher cat#EF0651). The D230A mutant of the GsSir2 protein was produced by QuikChange Site-Directed Mutagenesis 32 using respective mutagenic oligonucleotides (Supplementary Table 2). To generate the GsAgo protein containing a bulky His 6-StreppII-His 6-tag (HSH-tag, 29 aa.: LEGHHHHHSSWSHPQFEKGVEGHHHHHH) a whole operon of the GsSir2/Ago system was amplified by PCR from the genomic DNA using the oligonucleotides MZ-325 and MZ-326 (Supplementary Table 2), respectively. The resulting DNA fragment was digested by Eco31I and XhoI cloned into pBAD24 vector through NcoI/XhoI sites to generate pBAD24-HSH expression vector. The GsSir2/Ago mutants (GsSir22APZ/Ago, GsSir2/Ago3MID and GsSir2/Ago3PIW1) of the putative surface of the interaction with nucleic acids were designed based on the GsSir2/Ago structural model (see, Supplementary methods) changing positively charged residues that are structurally equivalent to the RsAgo residues involved in the interaction with the guide and the target to negatively charged residues in the corresponding domains (APAZ: R440E/R442E/R517E/R525E/R528E/R531E, MID: K155E/H166E/K170E/K207E, PIWI: R269E/K270E/H310E/K441E). The surface mutants were obtained using whole gene synthesis and cloning service provided by Twist Bioscience.

A PgSir2-Ago gene (BgramDRAFT_6510, WP_052303232.1) was amplified by PCR from the genomic DNA of Paraburkholderia graminis C4D1M (ATCC No. 700544) purchased from LGC Standards (UK) using the oligonucleotides MZ-915 and MZ-916 (Supplementary Table 2), respectively. The resulting DNA fragment was digested by BveI (ThermoFisher cat#FD1744) and HindIII (ThermoFisher cat#FD0504) and ligated into pBAD24 expression vector precleaved with NcoI and HindIII and dephosphorylated using FastAP.

The E. coli codon optimized genes (IDT Codon Optimization Tool) encoding the Sir2 (WP_053571900.1) and Ago (WP_053571899.1) of the CcSir2/Ago system (from Caballeronia cordobensis, NCBI taxon_id 1353886) were synthesized and cloned into pBAD/HisA expression vector by Twist Bioscience. The CcSir2 protein contains at its N-terminus a His-tag that can be cleaved by TEV protease. For purification of the CcSir2/Ago complex, a TwinStrep-tag (35 aa.: MGSAWSHPQFEKGGSGGSGGSAWSHPQFEKGS) was additionally fused to the N-terminus of the CcSir2 protein already containing a His-tag.

To swap ori regions between pCOLA and pCDF plasmids the DNA fragments containing ColA and CloDF13 ori were amplified by PCR using the oligonucleotides MZ-1217/MZ-1218 and MZ-1230/MZ-123, respectively (Supplementary Table 2). The resulting DNA fragments were digested by NheI and XbaI (ThermoFisher cat#FD0684) and ligated into pCOLA and pCDF vectors. precleaved with NheI and XbaI and dephosphorylated using FastAP.

To swap streptomycin resistance for kanamycin in the pCDF plasmid, plasmids pCDF and pCOLA were cleaved with NheI and Eco81I (ThermoFisher cat#FD0374) and isolated using a runView electrophoresis system (Cleaver Scientific). The purified fragments were then ligated into pCDF to yield a pCDF_Kn plasmid.

All gene sequences were confirmed by sequencing; links to DNA and protein sequences are presented in Supplementary Table 1.

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Phage restriction assay. E. coli MG1655 (ATCC 47076) cells carrying pBAD plasmids expressing wt or mutated Sir2/Ago systems were used for phage infection assays as described below. Whole-plasmid sequencing was applied to all transformed E. coli clones to verify the integrity of the system and lack of mutations, as

described before 33.
E. coli phages (T4, T7, lambda-vir) were kindly provided by U. Qimron. Phages SECphi17, SECphi18 and SECphi27 were isolated by the Sorek lab as described before 34. Small drop plaque assay was performed as described by Mazzocco et al. 35. Overnight culture of E. coli bacteria was diluted 1:100 in MMB medium (LB + 0.1 mM MnCl_2 + 5 mM MgCl_2 + 5 mM CaCl_2) supplied with 0.1% L-arabinose for expression induction. Bacterial cultures were incubated at 37 ° C until early log phase (OD_{600} = 0.3), and 500 l of bacteria were mixed with 25 ml of MMB agar (LB + 0.1 mM MnCl_2 + 5 mM MgCl_2 + 5 mM CaCl_2 + 0.5% agar + 0.1% L-arabinose) and poured into the square Petri dish. Serial dilutions of phage lysate in MMB were dropped on top of cell lain. After the drops dried up, plates were incubated at room temperature for 24 hours. Efficiency of plating (EOP) was determined via comparison of phage lysate titer on control bacteria and bacteria containing the Argonaute system with and without induction with L-arabinose.
Liquid culture phage infection experiments were performed as described previously by Ofir et al.33. After overnight incubation, the liquid suspension of pAgo-lacking and pAgo-containing E. coli cells were diluted 1:100 in MMB medium supplied with 0.2% L-arabinose and dispensed into a 96-well plate by 180-l volume. Plates were incubated at 37 ° C until the early log phase (OD600 = 0.3), then 20 l of phage lysate was added to each well in multiplicity of infection 5, 0.5, or 0.05, and each experiment was performed in three replicates. Optical density measurements at a wavelength 600 nm were taken every 15 min using a TECAN Infinite 200 plate reader.
Plasmid interference assay. E. coli BL21-AI and DH10B strain cells were pre-transformed with pBAD/HisA plasmid encoding a GsSir2/Ago (either wild-type or mutant variants), CcSir2/Ago or PgSir2-Ago - under the control of araBAD promoter. After 2 hours of induction with either 0.01% (w/v) (CcSir2/Ago) or 0.1% (GsSir2/Ago and PgSir2-Ago) L-arabinose at 37 ° C 200 RPM, cells were heat-shock transformed with pCDF (pCDF_Kn in the case of PgSir2-Ago in DH10B), pCOLA, pACYC184 or pRSF plasmids. After recovery, cells were either serially diluted and aliquots spotted on selection medium or undiluted suspensions spread on selection medium and CFUs counted manually. In parallel, viability and over-expression of pAgo-containing E. coli cells were monitored using serial dilutions and Western blot, respectively.
In separate experiments, E. coli BL21-AI strain cells were heat-shock transformed with both GsSir2/Ago operon-encoding pBAD/HisA construct and pCDF plasmid. Protein expression in selected double transformants was then induced by the addition of L-Ara (final concentration of 0.1% (w/v)) into liquid LB culture. After a 2-hour induction at 37 ° C 200 RPM, optical density (OD600) equalized, cultures were serially diluted and aliquots were spotted on a selection medium containing different antibiotics.
84 Expression and purification of GsSir2/Ago complexes. For GsSir2/Ago protein expression E. coli DH10B strain was transformed with a respective plasmid (Supplementary Table 1). Cells were grown at 37 ° C in LB medium in the presence of 50 g/ml ampicillin until OD600 = 0.7 was reached. Then, the temperature was decreased to 16 ° C and proteins were expressed for 16 h by adding 0.2% w/v L-arabinose. Next, harvested cells were disrupted by sonication in buffer A (20 mM Tris－HCl (pH 8.0 at 25 ° C), 500 mM NaCl, 2 mM phenylmethylsulfonyl fluoride, 5 mM 2-mercaptoethanol), and cell debris was removed by centrifugation. GsSir2/Ago complexes were purified to >90% homogeneity by chromatography through HisTrap HP chelating, HiTrap Heparin HP and HiLoad Superdex 200 columns (GE Healthcare). Purified proteins were stored at −20 ° C in
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

a buffer containing 20 mM Tris-HCl (pH 8.0 at 25 ° C), 200 mM KCl, 1 mM DTT and 50% v/v glycerol. The identity of the purified proteins was confirmed by mass spectrometry. Protein concentrations were determined from OD280 measurements using the theoretical extinction coefficients calculated with the ProtParam tool available at http://web.expasy.org/protparam/. GsSir2/Ago complex concentrations are

expressed in terms of heterodimer. The GsSir2/AgoMID surface mutant could not be purified due to its poor expression.

Antibodies used in this work. For Western blot analysis of target proteins, the following antibodies were used: 6x-His Tag monoclonal antibody (ThermoFisher, cat. #MA1-21315, RRID AB_557403, Lot # WE323793, clone HIS.H8); Goat anti-Mouse IgG (H+L) Secondary Antibody, AP conjugated (ThermoFisher, cat. #31320, RRID AB_228304, Lot # VH311913, polyclonal).

SEC-(MALS) and mass photometry. Size-exclusion chromatography of GsSir2/Ago complexes was carried out at room temperature using Superdex 200 10/300 GL column (GE Healthcare) pre-equilibrated with a buffer (20 mM Tris-HCl (pH 8.0 at 25 ° C), 500 mM NaCl). A calibration curve was generated by measuring the elution volumes of a series of standard proteins of known molecular mass (Bio-Rad). The molecular masses of pAgos complexes were calculated by interpolating their elution volume onto the calibration curve. SEC-MALS of GsSir2/Ago and CcSir2/Ago complexes was performed at room temperature using Superdex 200 10/300 GL column (GE Healthcare) pre-equilibrated with a buffer (20 mM Tris-HCl (pH 8.0 at 25 ° C), 500 mM NaCl, 0.03% NaN_3, 1 mM DTT), at 0.4 ml/min flow rate. Sample concentrations were 6 M and 6.5 M for GsSir2/Ago and CcSir2/Ago, respectively. The light scattering signals were monitored on a miniDawn TREOS II detector, concentrations of protein samples were measured using an Optilab T-rEX refractive index detector (Wyatt Technologies). Data were analysed in Astra software (Wyatt Technologies) using dn/dc value of 0.185 mL/g. Mass photometry of the GsSir2/Ago complex was performed using a Refeyn OneMP system (Refeyn). The protein complex was diluted to 20 nM in a buffer containing 20 mM Tris-HCl pH 8.0, 500 mM NaCl before measurement.

Small-angle X-ray scattering (SAXS) analysis. The synchrotron SAXS data were collected at beamline P12 operated by EMBL Hamburg at the PETRA III storage ring (DESY, Hamburg, Germany)36. GsSir2/Ago sample in the storage buffer was transferred into the sample buffer (20 mM Tris-HCl pH7.5, 200 mM NaCl, 5 mM MgCl, 2 mM β-mercaptoethanol) using gelfiltration NAP column (GE Healthcare) and concentrated by ultrafiltration to 1.2, 1.3, 1.6 and 5.5 mg/ml concentrations. The data were collected at the wavelength of 0.124 nm and the distance to the detector (Pilatus 2M, Dectris) was set to 3 m. Samples in the sample changer were kept at 10 ° C, capillary temperature was set to 20 ° C. Twenty frames exposed 0.045 sec were averaged for each concentration. The s-range of collected data was from 0.0133796 to 3.7925 nm -1. The data were analysed using programs of ATSAS 2.8.4 (r10552) suite 37. Data were normalized to an absolute scale with water as standard. As the data collected for the sample with a concentration of 1.22 mg/ml were noisy at higher s (Extended Data Fig. 4), and higher concentration data showed more aggregation at low s, we used a merged dataset produced with PRIMUS 38. Scattering data were parameterized and indirectly Fourier transformed with GNOM 539. Structural parameters of this dataset are summarized in Supplementary Table 3. Dimensionless Kratky plot in Extended Data Fig. 4 was calculated as described previously 40. The ab initio models were calculated by GASBOR 41 software. Molecular mass estimations of apo GsSir2/Ago complex in solution assessed by ATSAS tools (DATVC, DATMW) and server SAXSMoW (http://saxs.ifsc.usp.br/)42 are presented in the Supplementary Table 3.

Nucleic acid binding assay. The oligonucleotide substrates (Supplementary Table 2) were 5'- labelled with [γ-32P]ATP (PerkinElmer) and T4 polynucleotidyl kinase (PNK) (ThermoFisher cat#EK0031). The 3'-labeled substrate was prepared with [α-32P]cordycepin-5'-triphosphate

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

(Hartmann Analytics) and terminal deoxynucleotidyl transferase. An aliquot of the 3'-labelled substrate was subsequently phosphorylated with cold ATP and T4 PNK to obtain a 3'[α-32P]cordycepin-labelled oligonucleotide containing 5'-phosphate. Annealing was performed in the PNK reaction buffer supplemented with 50 mM EDTA at 2 M total single-stranded oligonucleotide concentration. Circular ssDNA substrate was

prepared by circularisation of 5'-32P labelled TK-49 using CircLigase II (Lucigen cat#CL9021K), according to manufacturer recommendations, and purified from a denaturing PAA gel (21% 29:1 acrylamide/bis-acrylamide in TBE supplemented with 8 M urea) by phenol-chloroform extraction, precipitated in 96 % ethanol with 0.45 M sodium acetate, washed with 75% ethanol, and resuspended in water.
For EMSA experiments, appropriate substrates and proteins were pre-diluted to 2x the final binding reaction concentration in 40 mM Tris-acetate (pH 8.3 at 23 ° C), 1 mM EDTA (TAE, Invitrogen cat#24710-030), supplemented with 5 mM magnesium acetate, 0.1 mg/ml BSA, 1 mM DTT, and 10% glycerol. The binding reactions were conducted by mixing equal volumes of enzyme and radiolabelled substrate. In all cases, final binding reactions contained 0.1 nM of radiolabelled substrate at 0-2 nM (0; 0.02; 0.05; 0.1; 0.2; 0.5; 1; 2) or 0-500 nM (0; 5; 10; 20; 50; 100; 200; 500) of GsSir2/Ago and CcSir2/Ago complexes. Three independent replicates were performed. For clarity, only EMSA gels obtained using low protein concentrations (0-2 nM) were shown, while for calculations of high K_d values EMSA data obtained using high protein concentrations (0-500 nM) were used.
Binding experiments for the binary GsSir2/Ago complex were conducted by first pre-mixing 5'-phosphorylated ssRNA or ssDNA guide with the equimolar GsSir2/Ago complex in the same buffer as above. The binary GsSir2/Ago:NA guide complex was then diluted to 2x final reaction concentration (in respect to guide) in the same buffer and mixed with a complementary 5'-32P-target oligonucleotide in the presence or absence of 67 ng/l heparin sodium salt (SigmaAldrich cat#). The final reaction contained 10 pM target NA and 0, 0.02, 0.05, 0.1, 0.2 0.5, and 1 nM of GsSir2/Ago:NA guide complex. Control (Cg*) contained 0.1 nM GsSir2/Ago:NA guide complex with the guide labelled with [γ-32P]ATP and 10 pM unlabeled target NA. Three independent replicates were performed.
The binding reaction mixtures were analysed by electrophoresis mobility shift assay (EMSA) in a PAA gel (8% 29:1 acrylamide/bis-acrylamide in TAE). The electrophoresis TAE buffer was supplemented with 5 mM magnesium acetate. Radiolabelled substrates were detected and quantified using a phosphor imager. The results were analysed with OptiQuant and OriginPro software. The K_d was calculated from the following formula:

$$\text{473} \qquad S_{NB} = A1 + \frac{\frac{100}{S_0}\left(\left(S_0\frac{(100-A1)}{100}\right)-E_0-K_D+\sqrt{\left(S_0\cdot\left(\frac{100-A1}{100}\right)+E_0+K_D\right)^2-4S_0\cdot\left(\frac{100-A1}{100}\right)\cdot E_0}\right)}{2}$$

where S_{NB - unbound substrate, nM; S_0 - initial substrate concentration, nM; E_0 - initial protein complex concentration, nM; K_d - dissociation constant, Al - nonbinding fraction of substrate, %.
Nucleic acid extraction and analysis. To obtain GsSir2/Ago-bound nucleic acids, E. coli DH10B was transformed with pBAD/HisA_TwinStrep_TEV_GsSir2/Ago and pCDF_Kn plasmids (Supplementary Table 1). Cells were grown at 37 ° C in LB medium in the presence of 50 g/ml ampicillin and 25 g/ml kanamycin until OD_{600} = 0.7 was reached. Then, expression was induced by adding 0.1% w/v L-arabinose, and cells were harvested after 2 h. Cells were disrupted using B-PER Bacterial Protein Extraction Reagent (ThermoFisher cat#78248) containing 6 mg/ml
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

lysozyme. The GsSir2/Ago-NA complex was purified as described above, instead, all buffer solutions contained 100 mM NaCl.
To extract nucleic acids co-purified with the GsSir2/Ago complex, 800 L of Rotiphenol/chloroform/isoamyl alcohol (Carl-Roth cat#A156) was added to the 800 L of purified protein-NA fractions in 5PRIME Phase Lock Gel tubes (Quantabio cat#733-2477). The upper aqueous phase was isolated and 0.1 volumes of 1 M sodium acetate, 3 volumes of 100% ethanol and 10 L glycogen (ThermoFisher cat#RO561) were added. This mixture was vortexed briefly and incubated at -20 ° C for 20 hours. Samples were centrifuged for 20 min

and the supernatant was removed from the pellet. The pellet was washed with cold (-20 ° C) 70% ethanol. The pellets containing the copurified nucleic acids were dried for 20 min at room temperature, and pellets were resuspended in 30 L water (free of nucleases).

Co-purified nucleic acids were dephosphorylated with FastAP Thermosensitive Alkaline Phosphatase (ThermoFisher cat# EF0651) and [γ-32P]-ATP (PerkinElmer) labelled with T4 polynucleotide kinase (PNK) (ThermoFisher cat#EK0031). Labelled nucleic acids were incubated with nucleases (ThermoFisher DNase I cat#18047019, RNase A/T1 cat# EN0551) for 30 min at 37 ° C. After nuclease treatment, samples were mixed with RNA Gel Loading Dye (ThermoFisher cat# R0641), heated for 5 min at 95 ° C and resolved on 20% denaturing (8 M Urea) polyacrylamide gels. The molecular weight marker used for RNA size identification was Decade Marker System (Ambion cat#AM7778) and 22 nt long RNA oligonucleotide. Radioactivity was captured from gels using phosphor screens and imaged using a Typhoon FLA 7000 laser-scatter (GE Healthcare).

In a control sample, a total RNA from induced cells was extracted using SPLIT RNA Extraction Kit (Lexogen cat#008). Then rRNA was removed using RiboCop for Gram-negative Bacteria (Lexogen cat#126).

RNA sequencing and analysis. Half of extracted RNA was treated with T4 PNK (ThermoFisher cat#EK0031) according to the protocol of the manufacturer. Then T4 PNK treated and untreated RNA samples were converted to DNA libraries using Small RNA-Seq Library Prep Kit (Lexogen cat#052). Concentration and quality of libraries were measured with Qubit Fluorometer (ThermoFisher) and 2100 Bioanalyzer (Agilent). Both libraries were sequenced using Illumina MiniSeq sequencing with single-end reads and 75 bp read length. Single-end reads were processed by trimming adapters with AdapterRemoval v2.3.045. Then the processed reads were aligned to the E. coli str. K12 strain DH10B genome (GenBank: CP000948.1) and the additional pBAD/HisA_TwinStrep_TEV_GsSir2/Ago, pCDF_Kn plasmids (Supplementary Table 1) using BWA-MEM v0.7.1744. In order not to filter out shorter reads during the alignment process, aligned reads with MAPQ values greater or equal to 15 were chosen. FastQC v0.11.845 was used for read quality control and SAMtools v1.746 - for indexing, sorting and analysing alignment files. A custom script (fragmentation-bias.jl) in combination with Weblogo v3.7.447 were used to produce nucleotide frequency plots. The custom script had to be implemented to ensure that only aligned reads would be used for nucleotide frequency analysis. Gene enrichment analysis was performed with bedtools v2.26.048 and FPKM_count.py v4.0.0 of RSeqQC package49. IGV v2.5.250 was mainly used to inspect and visualise read coverage along the genomes. Control DNA library of a total RNA was prepared using CORALL Total RNA-Seq Library Prep Kit (Lexogen cat#095). Concentration and quality of the library was measured with Qubit Fluorometer (ThermoFisher) and 2100 Bioanalyzer (Agilent) according to the protocol of the manufacturer.

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Control DNA library was sequenced using Illumina NextSeq sequencing with pair-end reads and 75 bp read length. The read processing, alignment and alignment analysis were analogous to those samples from Illumina MiniSeq sequencing.

Preparation of E. coli cells for NAD + quantification. Overnight cultures of single colonies of E. coli DH10B strain harbouring a pBAD-His construct with either wt GsSir2/Ago or mutant system (GsSir2/Ago-HSH or GsSir2(D230A)/Ago), or empty vector (negative control) were diluted and grown in LB broth (BD) supplemented with respective antibiotics (50 g/ml ampicillin and 25 g/ml streptomycin) at 37 ° C until they reached OD600 of 0.4 - 0.5. Cell cultures were either induced to express the protein or not (control samples). L-Ara (0.1% final concentration) was added to induce protein expression. Induced and non-induced cultures were harvested 2 hours later. The cultures were normalized to OD600 of approximately 0.7 and the pellet from 1 mL of culture suspension was stored at -80 ° C until further analysis. All cell pellets were lysed by adding B-PER solution (Thermofisher Scientific) supplemented with 6 mg/ml lysozyme (62971, Fluka)

for 20 min at room temperature while gently rocking (Multi Bio 3D Mini-Shaker, Biosan). Cell debris was removed by centrifugation and metabolites were isolated by phenol:chloroform:isoamyl alcohol (PCI) (25:24:1, v/v/v) extraction. Metabolites were stored at -20 ° C until MS-HPLC analysis. Additionally, the endogenous NAD + concentration was estimated using NAD/NADH Quantitation Kit (Sigma Aldrich, cat# MAK037) from four independent measurements.

In vitro NADase assay. Reaction mixtures with a volume of 25 L were prepared with the following final concentrations: 0.5 M GsSir2/Ago or mutant complex, 50 M NAD+, 1× Tango buffer (33 mM Tris-acetate (pH 7.9 at 37 ° C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA, ThermoScientific, cat#BY5), 1 mM DTT, 0.5 M 5'P-RNA guide (TF-A) and/or 0.5 M ssDNA (MZ-949 or MZ-589) (Supplementary Table 2). Reactions with RNA guide were preincubated for 15 min at 37 ° C, then ssDNA was added and the mixture was incubated for 1 h at 37 ° C. 3 L of each sample was used as input for the NAD/NADH Quantitation Kit (Sigma-Aldrich, cat# MAK037) according to the instructions provided by the manufacturer. All experiments were performed in triplicates. These samples were also used for mass spectrometry.

Mass spectrometry of NAD +. To quantitate NAD + bound to GsSir2/Ago and CcSir2/Ago complexes high-performance liquid chromatography-mass spectrometry/mass spectrometry (HPLC-MS/MS) analysis was used. First, purified pAgos complexes were diluted to 5 M in a buffer containing 20 mM Tris-HCl (pH 8.0 at 25 ° C), 200 mM NaCl. Then 20 l of the solution was incubated at 70 ° C for 20 min and centrifuged for 30 min (16,100 g at 4 ° C) to remove unfolded proteins. The supernatants and NAD + standards were analysed by Electrospray Ionization mass spectrometry (ESI-MS) using an integrated HPLC/ESI-MS system (1290 Infinity, Agilent Technologies/Triple Quadrupole 6410, Agilent Technologies), equipped with a Supelco Discovery HS C18 column (7.5 cm × 2.1 mm, 3 m), Agilent Technologies. HPLC/ESI-MS/MS was performed using two ion transitions to detect NAD + in the samples: 662.1 → 540.1 and 662.1 → 426.0. Ion transition 662.1 → 540.1, as it is the most abundant, was used for the quantitative analysis. Mobile phase A was 5 mM ammonium acetate in water, pH 7.0 and mobile phase B was 5 mM ammonium acetate in methanol, pH 7.0. The HPLC parameters were as follows: flow 0.25 ml/min; column temperature 30° C; 0-3 min, 0%B; 3-9 min, 0-40%B; 9-10 min, 40-100%B; 10-13 min, 100%B. The MS was operated using negative electrospray ionisation at 2500 V, the gas temperature was set to 300 ° C, the fragmentor voltage was 135 V. Multiple reaction monitoring (MRM) was used with a collision energy of 15 V to measure ion m/z 540.1 (ion transition 662.1 → 540.1) and also with a collision energy of 20 V to measure ion m/z 426.0) (ion transition 662.1 → 426.0).

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

To quantitate endogenous NAD + HPLC-MS analysis was performed by Electrospray Ionization mass spectrometry (ESI-MS) using an integrated HPLC/ESI-MS system (1290 Infinity, Agilent Technologies/Q-TOF 6520, Agilent Technologies), equipped with a Supelco Discovery HS C18 column (7.5 cm × 2.1 mm, 3 m), Agilent Technologies. The samples were investigated in both negative and positive ionization modes. For negative ionization mode, solvents A (5 mM ammonium acetate in water pH 7.0) and B (5 mM ammonium acetate in methanol, pH 7.0) were used. For positive ionization mode, solvents C (0,02% formic acid in water) and D (0,02% formic acid in acetonitrile) were used. In both cases elution was performed with a linear gradient of solvents at a flow rate of 0.3 ml/min at 30 ° C as follows: 0 - 5 min, 0% B; 5 - 18 min, 20% B; 18 - 22 min, 100% B, 22 - 27 min 100% B. Ionization capillary voltage was set to 2500 V and fragmentor to 150V. A list of compounds that could be expected to be products of NAD + hydrolysis and relative m/z value is as follows: ADRPR [M-H] - m/z=558.0644, cADPR [M-H] - m/z =540.0538, AMP [M-H] - m/z=346.0558, cAMP [M-H] - m/z=328.0452, ADP [M-H] - m/z=426.0221, cADP [M-H] - m/z=408.0116, nicotinamide [M+H] + m/z = 123.0553, adenine [M+H] + m/z = 136.0618. Only traces of AMP and ADP were detected in all samples; other products from the list were absent.

NAD+ hydrolysis products generated by GsSir2/Ago in vitro were analysed as above. Using negative ionization mode only accumulation of ADPR was detected.

588 Data availability

All data are available in the paper and the supplementary material. In addition, small and total RNA sequencing data are available on the NCBI Sequence Read Archive under BioProject ID PRJNA851009. SAXS data are available in the Small Angle Scattering Biological Data Bank SASDBB under SASDBD ID SASDNH2: https://www.sasdb.org/project/1486/cuadir1lvf/. Plasmid sequences used in this work are available at https://www.benchling.com, with exact links for each plasmid provided in Supplementary Table 1. Geobacter sulfurreducens, Caballeronia cordobensis, and Paraburkholderia graminis genomes (respective GenBank accessions: GCA_000210155.1, GCA_001544575.2 and GCA_000172415.1) and all associated sequence and annotation data were obtained from NCBI (ftp://ftp.ncbi.nlm.nih.gov/genomes/Bacteria/). Searches through Pfam (http://pfam.xfam.org/), SwissProt (https://www.expasy.org/resources/uniprotkb-swiss-prot), and PDB (https://www.rcsb.org/) databases were performed. PDB structures mentioned in this study: 5AWH, 4N41, 5UX0, 6LHX, 2H4F.

602 Code availability

The Julia script used to identify nucleotide frequency in the beginning of the aligned reads and prepare input for Weblogo program is available at GitHub repository: https://github.com/agrybauskas/argonaute-bound-rna-manuscript

606 Acknowledgements

We thank the Siksnys laboratory members for their comments on the manuscript and fruitful discussion. This work was supported by European Social Fund [09.3.3-LMT-K-712-01-0126 to V.S.] under a grant agreement with the Research Council of Lithuania (LMTLT), the Israel Science Foundation [grant ISF 296/21 to R.S.], and the Deutsche Forschungsgemeinschaft [SPP 2330, grant 464312965]. The authors thank Tomás de Garay at Refeyn Ltd. for the mass photometry experiments. Funding for open access charge: Vilnius University.

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

613 Author Contributions

V.S. and M.Z. designed the study; K.T. and .V. performed bioinformatics and structural modelling; A.L. performed the phage restriction experiments; E.S., D.D., E.G., S.A., U.T. performed the plasmid transformation experiments; A.S. purified the proteins and performed the SEC-(MALS) experiments; R.G. performed the SEC experiments; E.M. performed the SAXS experiments; E.Z., E.G., D.D. and R.G. performed the EMSA experiments; E.Z. and E.J. reconstituted the GsSir2/Ago-RNA complex and performed the biochemical analysis; A.G. performed the RNA-seq analysis; A.R. performed the mass spectrometry analysis; D.D. performed the NAD + determination experiments in vivo; E.Z. performed the NAD + hydrolysis experiments in vitro; R.S., V.S. and M.Z. analysed the data; M.Z. wrote the initial manuscript with input from E.G., R.S., V.S. and other authors. All authors approved the final version.

624 Competing Interests statement

625 VS is the chairman of CasZyme. R.S. is the scientific founder of BiomX and Ecophage.

626 The remaining authors declare no competing interests.

627 Notes

During the revision of this manuscript, a paper was published that shows plasmid-induced degradation of NAD + in vivo by short pAgo-associated TIR-APAZ systems (named SPARTA) 52. The paper also shows that expression of Sir2/Ago systems (named SPARSA) in E. coli triggers NAD depletion.

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND

4.0 International license.

633 Tables
Table 1. Nucleic acid binding by Sir2/Ago and their binary complexes pre-loaded with either ssRNA or ssDNA guides. K_d values (averages ± standard deviation of three independent replicates) were calculated from EMSA data. Some binding experiments were performed in the presence of heparin (indicated).
Nucleic acid binding by Sir2/Ago			
Nucleic acid	Nucleic acid	Kd, nM	Kd, nM
Nucleic acid	Nucleic acid	GsSir2/Ago	CcSir2/Ago
ssDNA	ssDNA	0.033 ± 0.016	0.138 ± 0.007
ssRNA	ssRNA	0.042 ± 0.015	0.36 ± 0.098
circular ssDNA	circular ssDNA	0.072 ± 0.018	0.18 ± 0.068
dsRNA/DNA	dsRNA/DNA	4.31 ± 0.37	2.4 ± 0.69
dsDNA	dsDNA	51 ± 20	330 ± 87
dsRNA	dsRNA	73 ± 29	81 ± 15
Nucleic acid binding by GsSir2/Ago-gRNA and GsSir2/Ago-gDNA complexes			
Guide	Target	Kd, nM	Kd, nM
Guide	Target	- Heparin	+ Heparin
ssRNA	ssDNA	0.020 ± 0.008	0.079 ± 0.007
ssRNA	ssRNA	0.205 ± 0.058	0.567 ± 0.112
ssDNA	ssDNA	0.022 ± 0.005	n.b.*
ssDNA	ssRNA	0.252 ± 0.114	n.b.*
ssRNA	nsp-ssDNA#	0.420 ± 0.231	4.29 ± 2.11
ssDNA binding by GsSir2/Ago-gRNA variants			
Mutant	Mutant	Kd, nM	Kd, nM
Mutant	Mutant	- Heparin	+ Heparin
GsSir2(D230A)/Ago	GsSir2(D230A)/Ago	0.038 ± 0.013	0.140 ± 0.057
GsSir2/Ago-HSH	GsSir2/Ago-HSH	0.140 ± 0.017	0.203 ± 0.058
GsSir2APAZ/Ago	GsSir2APAZ/Ago	n.b.*	n.b.*
GsSir2/AgoPIWI	GsSir2/AgoPIWI	n.b.*	n.b.*
* No binding was detected under experimental conditions used. # A non-complementary ssDNA was used.			
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

639 Figures
A
Long pAgos
GsSir2/Ago from Geobacter sulfurreducens
CcSir2/Ago from Caballeronia cordobensis
Sir2	APAZ			MID	PIWI	
1	314	580	38	213	474	
				Catalytically inactive		
PgSir2-Ago from Paraburkholderia graminis
B
640
C
D

E
F
Fig. 1. Sir2/Ago provide defence against phages. A, Schematic representation of the subunit/domain composition of different pAgo variants. Catalytically active pAgos contain a conserved catalytic DEDX tetrad that is mutated in inactive pAgos. Catalytically inactive short pAgos lack the canonical PAZ domain, however, an accessory APAZ domain is present in putative Sir2, TIR or Mrr proteins associated with short pAgos. MID indicates middle; L - linker domain; N - N-terminal domain. Short pAgos from G. sulfurreducens, C. cordobensis, and P. graminis associated with Sir2 protein were studied in this work. B, Schematic diagram of phage restriction assays. C, Efficiency of plating (EOP) of lambda-vir and SECphi27 phages infecting E. coli cells with and without the GSir2/Ago system. The bar graphs show the number of p.f.u. as arithmetic means of two replicates in the absence and the presence of the inducer Larabinose (L-Ara), with individual data points superimposed. Grey bars represent EOP on pAgo-lacking cells and black bars - in pAgo-containing cells. Representative images of plaque assays are also presented. D, EOP of lambda-vir and SECphi27 phages infecting the wt and mutant GSir2/Ago systems in the presence of L-Ara. GSir2(D230A)/Ago and GSir2/Ago-HSH are variants that contain D230A mutation in the Sir2 domain or (HSH-tag) on the C-terminus of pAgo, respectively. The bar graphs show the number of p.f.u. as arithmetic means of three replicates, with individual data points superimposed. Grey bars represent EOP on pAgo-lacking cells and black bars - in pAgo-containing cells. E, EOP of lambda-vir and SECphi27 phages infecting pAgo-lacking cells and cells containing GSir2/Ago, CcSir2/Ago and PgSir2-Ago in the presence of L-Ara. The bar graphs show the number of p.f.u. as arithmetic means of three replicates, with individual data points superimposed. Grey bars represent EOP on pAgo-lacking cells and black bars - in pAgo-containing cells. F, Lambda phage infection in liquid cultures of E. coli cells containing the GSir2/Ago system. GSir2/Ago-lacking (shown in grey) or GSir2/Ago-containing (shown in orange) E. coli were infected at t=0 at multiplicities of infection (MOI) of 0.05, 0.5 and 5. Each curve represents one individual replicate; two replicates for each MOI are shown.
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
suites
18
A
B
C
D
E
Fig. 2. The GsSir2/Ago system interferes with plasmid transformation. A, Schematic representation of the experiment. B, Qualitative evaluation of plasmid transformation efficiency in E. coli cells carrying GsSir2/Ago system.
Top: comparison of cell viability in the presence or absence of plasmid-borne GsSir2/Ago expression.
Bottom: comparison of plasmid transformation efficiencies in the presence or absence of plasmid-borne GsSir2/Ago expression.
C, Left - comparison of pCDF transformation efficiency between cells expressing wt and mutant GsSir2/Ago complexes; right - quantification of transformation efficiencies (three independent replicates, the red line represents average transformation efficiency). D, Top - schematic representation of ori exchange between pCDF and pCOLA plasmids; bottom - comparison of plasmid transformation efficiencies. pCDF, pCDF with CloDF13 ori exchanged with ColA ori (pCDF_ColA), pCOLA and pCOLA with ColA ori exchanged with CloDF13 ori (pCOLA_CloDF13) plasmids were used for transformation of E. coli cells carrying GsSir2/Ago system. E, Cell viability in the absence of antibiotic selection. In the case of the wt GsSir2/Ago system, the cell

viability decreases on the plates even in the absence of Cb and Str antibiotics suggesting that GsSir2/Ago in the presence of the pCDF plasmid triggers cell death.
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
19
A
B
C
D
677
Fig. 3. The GsSir2 and GsAgo proteins form a heterodimeric complex. A, Top - SDS PAGE of purified wt GsSir2/Ago, a D230A mutant, and C-terminal HSH tag-bearing GsSir2/Ago. Red star in the HSH-tagged sample lane marks an overlap of bands in the gel due to similar mass. Bottom - anti-His-tag Western blot of same samples. The red star shows the lane where the His-tag is on the C-terminus of Ago, rather than the N-terminus of Sir2. Two replicates. B, Mass photometry data of the GsSir2/Ago complex, with masses and respective particle population counts indicated. According to mass spectrometry of the purified GsSir2/Ago complex (Extended Data Fig. 3F), the molar mass of the GsSir2/Ago heterodimer is 121 kDa. C, SEC-MALS data of GsSir2/Ago, showing the chromatography peak and molar weight of the Sir2/Ago heterodimer. D, Semitransparent space-filling ab initio model of GsSir2/Ago calculated from SAXS data with a fitted-in AF GsSir2/Ago model in cartoon representation. Colour coding: Sir2 domain - cyan, APAZ - brown, MID - yellow, PIWI - green.
688
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
20
A
B
E
689
Fig. 4. Nucleic acid binding by GsSir2/Ago in vitro and in vivo. A, Binding of RNA or DNA targets by GsSir2/Ago binary complexes pre-loaded with 5'P-RNA or 5'P-DNA guides. In EMSA experiments the pre-formed GsSir2/Ago-gNA binary complex was mixed with a radiolabelled target strand indicated by the asterisk (see “Materials and Methods” for the details). To show that no displacement of the guide by the target strand occurs under these experimental conditions, a control (Cg) experiment was performed where the guide, rather than the target, was radioactively labelled. Only the pre-annealed RNA/DNA heteroduplex was loaded in the control lane “d”. Three independent replicates performed. B, GsSir2/Ago co-purifies with small RNAs. Nucleic acids that co-purified with GsSir2/Ago were, first, dephosphorylated, then, [γ-32P]-ATP radiolabelled and treated with DNase I or RNase A/T1, or both, and resolved on a denaturing polyacrylamide gel. R, control 22 nt RNA oligonucleotide; M, RNA ladder Decade Marker System (Ambion). Three independent replicates performed. C, Length distribution of small RNA co-purified with GsSir2/Ago as determined by sequencing. “5'P-RNA” sample means that only small RNAs containing 5'-phosphate were sequenced, while in “5'P+5'OH-RNA” sample both 5'-phosphate or 5'-OH bearing small RNAs were sequenced. D, Small RNAs associated with GsSir2/Ago show 5'-AU preference. E, Distribution of small RNAs co-purified with GsSir2/Ago from the E. coli host carrying the pCDF_Kn target and pBAD_GsSir2/Ago expression plasmids (above). IGV viewer representation of total RNA extracted from E. coli is shown below. Cartoons indicate

promoters (P_BAD), protein-coding genes (KanR, LacI, GsSir2, GsAgo, AmpR, AraC), plasmid ori (CloDF13, ColE1) and their RNAI and RNAII transcripts. Red arrow shows RNase H cleavage site in RNAII required for initiation of DNA strand synthesis during plasmid replication.
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
21
A
B
C
D
E
F
Fig. 5. GsSir2/Ago binds and hydrolyses NAD+. A, the D230A mutation within the Sir2 protein abolished NAD+ binding. B, NAD+ amounts in E. coli cells in the presence of the (non)induced wt and mutant GsSir2/Ago systems and in the presence and absence of the pCDF plasmids. NAD+ amounts were estimated according to the EIC areas of NAD+ (m/z 662.1018). pBAD – empty vector; wt – GsSir2/Ago; D230A – GsSir2(D230A)/Ago; HSH – GsSir2/Ago-HSH. C, NAD+ depletion by GsSir2/Ago in vitro. The wt GsSir2/Ago or mutant complex (0.5 M) was incubated with NAD+ (50 M) for 1 h at 37 ° C (see experimental details in “Materials and Methods”). gRNA, 5’P-RNA guide; tDNA, target DNA complementary to the RNA guide; nsp-DNA, ssDNA non-complementary to the RNA guide. D, NAD+ hydrolysis by wt GsSir2/Ago in vitro. The binary GsSir2/Ago-gRNA complex (0.5 M) was incubated with NAD+ (50 M) for 1 h at 37 ° C in the presence of the complementary DNA target (0.5 M). NAD+ depletion and ADPR accumulation was analysed by MS according to the EIS of NAD+ (m/z 662.1018) and ADPR (m/z 558.0644), respectively. AC, the activated wt GsSir2/Ago-gRNA/tDNA complex. E, Representative mass chromatograms of NAD+ hydrolysis by wt GsSir2/Ago in vitro (as in D). C, a control sample without GsSir2/Ago; AC, the activated wt GsSir2/Ago-gRNA/tDNA complex. F, Putative model of GsSir2/Ago defence against mobile genetic elements. After lambda phage infection or pCDF plasmid transformation, GsSir2/Ago acquires small 5’-AU-RNAs originated from the invader transcripts (e.g., from pCDF ori region). The GsSir2/Ago binary complex, guided by small RNA, targets the invaders’ complementary DNA, becomes activated and hydrolyses NAD+ resulting in a cell death.
726
727
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
22
728 References
1. Kuhn, C. D. & Joshua-Tor, L. Eukaryotic Argonautes come into focus. Trends Biochem. Sci. 38, 263–271 (2013).
2. Pratt, A. J. & MacRae, I. J. The RNA-induced silencing complex: A versatile gene-silencing machine. J. Biol. Chem. 284, 17897–17901 (2009).
3. Sheu-Gruttadauria, J. & MacRae, I. J. Structural Foundations of RNA Silencing by Argonaute. J. Mol. Biol. 429, 2619–2639 (2017).
4. Olina, A. V., Kulbachinskiy, A. V., Aravin, A. A. & Esyunina, D. M. Argonaute Proteins and Mechanisms of RNA Interference in Eukaryotes and Prokaryotes. Biochem. 83, 483–497 (2018).
5. Hutvagner, G. & Simard, M. J. Argonaute proteins: Key players in RNA silencing. Nat. Rev. Mol. Cell Biol. 9, 22–32 (2008).

6. Swarts, D. C. et al. The evolutionary journey of Argonaute proteins. Nat. Struct. Mol. Biol. 21, 743–753 (2014).
7. Kwak, P. B. & Tomari, Y. The N domain of Argonaute drives duplex unwinding during RISC assembly. Nat. Struct. Mol. Biol. 19, 145–151 (2012).
8. Ryazansky, S., Kulbachinskiy, A. & Aravin, A. A. The Expanded Universe of Prokaryotic Argonaute Proteins. MBio 9, 1–20 (2018).
9. Hegge, J. W., Swarts, D. C. & Van Der Oost, J. Prokaryotic argonaute proteins: Novel genome-editing tools? Nat. Rev. Microbiol. 16, 5–11 (2018).
10. Lisitskaya, L., Aravin, A. A. & Kulbachinskiy, A. DNA interference and beyond: structure and functions of prokaryotic Argonaute proteins. Nat. Commun. 9, 1–12 (2018).
11. Willkomm, S., Makarova, K. S. & Grohmann, D. DNA silencing by prokaryotic Argonaute proteins adds a new layer of defense against invading nucleic acids. FEMS Microbiol. Rev. 42, 376–387 (2018).
12. Kuzmenko, A. et al. DNA targeting and interference by a bacterial Argonaute nuclease. Nature (2020) doi:10.1038/s41586-020-2605-1.
13. Swarts, D. C. et al. DNA-guided DNA interference by a prokaryotic Argonaute. Nature 507, 258–261 (2014).
Jolly, S. M. et al. Thermus thermophilus Argonaute Functions in the Completion of DNA Replication. Cell 182, 1545-1559.e18 (2020).
15. Olovnikov, I., Chan, K., Sachidanandam, R., Newman, D. K. & Aravin, A. A. Bacterial Argonaute Samples the Transcriptome to Identify Foreign DNA. Mol. Cell 51, 594–605 (2013).
16. Liu, Y. et al. A programmable omnipotent Argonaute nuclease from mesophilic bacteria. Kurthia massiliensis. Nucleic Acids Res. 1–12 (2021) doi:10.1093/nar/gkaa1278.
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

17. Kropocheva, E., Kuzmenko, A., Aravin, A. A., Esyunina, D. & Kulbachinsky, A. A programmable pAgo nuclease with universal guide and target specificity from the mesophilic bacterium Kurthia massiliensis. Nucleic Acids Res. 1, 1–28 (2021).
18. Makarova, K. S., Wolf, Y. I., van der Oost, J. & Koonin, E. V. Prokaryotic homologs of Argonaute proteins are predicted to function as key components of a novel system of defense against mobile genetic elements. Biol. Direct 4, 29 (2009).
19. Burroughs, A. M., Ando, Y. & Aravind, L. New perspectives on the diversification of the RNA interference system: Insights from comparative genomics and small RNA sequencing. Wiley Interdiscip. Rev. RNA 5, 141–181 (2014).
North, B. J. & Verdin, E. Protein family review Sirtuins: Sir2-related NAD-dependent protein deacetylases Gene organization and evolutionary history. Genome Biol. 5, 224 (2004).
21. Gallego-Jara, J. et al. Bacterial Sirtuins Overview: An Open Niche to Explore. Frontiers in Microbiology vol. 12 (2021).
22. Wang, Y., Sheng, G., Juranek, S., Tuschl, T. & Patel, D. J. Structure of the guide-strand-containing argonaute silencing complex. Nature 456, 209–213 (2008).
23. Frank, F., Sonenberg, N. & Nagar, B. Structural basis for 5'-nucleotide base-specific recognition of guide RNA by human AGO2. Nature 465, 818–822 (2010).
24. Ka, D., Oh, H., Park, E., Kim, J. H. & Bae, E. Structural and functional evidence of bacterial antiphage protection by Thoiris defense system via NAD+ degradation. Nat. Commun. 11, 1–8 (2020).
Kim, S., Jung, Y. & Lim, D. Argonaute system of Kordia jejudonensis is a heterodimeric nucleic acid-guided nuclease. Biochem. Biophys. Res. Commun. 525, 755–758 (2020).
26. Dasgupta, S., Masukata, H. & Tomizawa, J. ichi. Multiple mechanisms for initiation of ColE1 DNA

replication: DNA synthesis in the presence and absence of ribonuclease H. Cell 51, 1113–1122 (1987).
27. del Solar, G., Giraldo, R., Ruiz-Echevarria, M. J., Espinosa, M. & Diaz-Orejas, R. Replication and Control of Circular Bacterial Plasmids. Microbiol. Mol. Biol. Rev. 62, 434-464 (1998).
28. Selzer, G., Som, T., Itoh, T. & Tomizawa, J. ichi. The origin of replication of plasmid p15A and comparative studies on the nucleotide sequences around the origin of related plasmids. Cell 32, 119–129 (1983).
29. Garb, J. et al. Multiple phage resistance systems inhibit infection via SIR2-dependent NAD+depletion. bioRxiv 2021.12.14.472415 (2021) doi:10.1101/2021.12.14.472415.
30. Tal, N. et al. Cyclic COMP and cyclic UMP mediate bacterial immunity against phages. Cell 184, 5728-5739.e16 (2021).
31. Ofir, G. et al. Antiviral activity of bacterial TIR domains via immune signalling molecules. Nature 600, 116–120 (2021).
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

32. Zheng, L., Baumann, U. & Raymond, J. L. An efficient one-step site-directed and site-saturation mutagenesis protocol. Nucleic Acids Res. 32, (2004).
33. Ofir, G. et al. DISARM is a widespread bacterial defence system with broad anti-phage activities. Nat. Microbiol. 3, 90–98 (2018).
34. Doron, S. et al. Systematic discovery of antiphage defense systems in the microbial pangenome. Science (80-.). 359, ear4120 (2018).
35. Mazzocco, A., Waddell, T. E., Lingohr, E. & Johnson, R. P. Enumeration of Bacteriophages Using the Small Drop Plaque Assay System. Methods Mol. Biol. 81–85 (2009) doi:10.1007/978-1-60327-164-6_9.
36. Blanchet, C. E. et al. Versatile sample environments and automation for biological solution X-ray scattering experiments at the P12 beamline (PETRA III, DESY). J. Appl. Crystallogr. 48, 431–443 (2015).
37. Franke, D., Petoukhov, M. V, Konarev, P. V & Panjkovich, A. ATSAS 2.8: a comprehensive data analysis suite for small-angle scattering from macromolecular solutions. J. Appl. Crystallogr. 1212–1225 (2017) doi:10.1107/S1600576717007786.
38. Konarev, P. V., Volkov, V. V., Sokolova, A. V., Koch, M. H. J. & Svergun, D. I. PRIMUS - a Windows-PC based system for small-angle scattering data analysis. J. Appl. Crystallogr. 36, 1277–1282 (2003).
Svergun, D. I. Determination of the regularization parameter in indirect-transform methods using perceptual criteria. J. Appl. Crystallogr. 25, 495–503 (1992).
Durand, D. et al. NADPH oxidase activator p67phox behaves in solution as a multidomain protein with semi-flexible linkers. J. Struct. Biol. 169, 45–53 (2010).
Svergun, D. I., Petoukhov, M. V. & Koch, M. H. J. Determination of domain structure of proteins from x-ray solution scattering. Biophys. J. 80, 2946–2953 (2001).
42. Fischer, H., De Oliveira Neto, M., Napolitano, H. B., Polikarpov, I. & Craievich, A. F. Determination of the molecular weight of proteins in solution from a single small-angle X-ray scattering measurement on a relative scale. J. Appl. Crystallogr. 43, 101–109 (2010).
43. Schubert, M., Lindgreen, S. & Orlando, L. AdapterRemoval v2: rapid adapter trimming, identification, and read merging. BMC Res. Notes 9, 88 (2016).
44. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25, 1754–1760 (2009).
45. Andrews, S. & others. FastQC: a quality control tool for high throughput sequence data. (2010).

46. Li, H. et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079 (2009).
47. Crooks, G. E., Hon, G., Chandonia, J.-M. & Brenner, S. E. WebLogo: A Sequence Logo Generator. Genome Res. 14, 1188–1190 (2004).
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

48. Quinlan, A. R. BEDTools: The Swiss-Army Tool for Genome Feature Analysis. Curr. Protoc. Bioinforma. 47, 11.12.1-11.12.34 (2014).
49. Wang, L., Wang, S. & Li, W. RSeQC: quality control of RNA-seq experiments. Bioinformatics 28, 2184–2185 (2012).
50. Thorvalsdottir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Brief. Bioinform. 14, 178–192 (2013).
51. Kaya, E. et al. A bacterial Argonaute with noncanonical guide RNA specificity. Proc. Natl. Acad. Sci. U. S. A. 113, 4057–4062 (2016).
52. Koopal, B. et al. Short prokaryotic Argonaute systems trigger cell death upon detection of invading DNA. Cell 185, 1–16 (2022).
850
851
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

852 Extended data figures
A
B
C
Sir2-APAZ group Ib
Sir2-APAZ group Ia
ThsAhomologues
Long-B
kU
kU
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Extended Data Fig. 1. Bioinformatic analysis. A, PIWI catalytic tetrad DEDX alignment. The 4 catalytic residues (red numbers indicate positions of corresponding GsAgo positions) are shown in 4 motifs of ±3 positions. The motifs are separated by vertical blue lines. Sequence names consist of the following: NCBI sequence ID, abbreviated phylum (e.g., 'ProG' - gamma-proteobacteria) and organism name. B, Top - a circular phylogenetic tree was generated according to supplementary data provided with Ryazansky et al.8 Long-A pAgo variants are coloured in green (truncated variants without the PAZ domain, light green), long-B pAgo proteins are light green (truncated variants without PAZ, green), and short pAgo proteins are orange. pAgo proteins containing the catalytic tetrad DEDX in their PIWI domain are indicated in blue on the outer circle; pAgos with inactivated PIWI domain are indicated in light grey on the outer circle. pAgo

proteins of the GsSir2/Ago, CcSir2/Ago and PgSir2-Ago systems are indicated by 'Gs', 'Cc' and 'Pg', respectively. Bottom - Circular phylogenetic tree of APAZ domains. The circular phylogenetic tree of the five groups of APAZ domains was generated using APAZ domain alignments from Ryazansky et al.8 supplementary file 7. C, Top - Combined alignment of Sir2 domains. Alignment consists of 3 parts, separated by horizontal black lines. In the top part, the Sir2 domain sequences of the GsSir2, CcSir2, PgSir2-Ago and homologues are shown. Logos above depict the conservation of Sir2 domains of Ia and Ib groups. The indicated position numbers correspond to the GsSir2 sequence. In the bottom part, homologues (sirtuins) of catalytically active Thermotoga maritima Sir2 (TmSir2) deacetylase are shown. Logos below indicate the conservation of these homologues. The position numbers correspond to the TmSir2 sequence. Sequences of six motifs that include all positions that form the NAD --binding pocket, as seen in the TmSir2 structure (PDB ID 2H4F) are shown. Sequence names for the top alignment consist of sequence ID, abbreviated phylum and organism name. Sequence names for bottom alignment all start with "Sir2hom" followed by sequence ID, organism name and short protein name (based on annotation). Stars above the logos indicate residues in the NAD --binding pocket of canonical sirtuins (e.g., TmSir2) that are also conserved. Star colours indicate conservation between the two groups: green - conserved in both canonical sirtuins and GsSir2-like; blue - conserved in both groups, but different; yellow - conserved only in canonical sirtuins; red - conserved only in GsSir2-like proteins. In the middle, alignment of ThsA homologues with Sir2 domains. Bottom - MID domain alignment. Red stars indicate positions of amino acids involved in the binding of the 5'-P end of the guide nucleic acid. The numbering above corresponds to the GsAgo sequence. Additionally, concatenated alignment of just the 6 indicated positions is shown on the right. The three sequences of interest are indicated with red rectangles. Numbers on the left and right of the alignment indicate the first and last positions in the alignment for each sequence.
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
28
A
B
C
D
G
F
the absence and in the presence of the inducer L-arabinose (L-Ara), with individual datapoints superimposed. Grey bars
887
represent efficiency of plating (EOP) on pAgo-lacking cells and black bars are EOP in pAgo-containing cells. B, Left -
888
qualitative characterization of plasmid restriction capabilities of PgSir2-Ago system in E. coli strain DH10B. Top:
889
comparison of cell viability in the presence or absence of plasmid-borne PgSir2-Ago expression. Bottom: comparison
890
of plasmid transformation efficiencies in the presence or absence of plasmid-borne PgSir2-Ago expression. Right -
891
qualitative characterization of plasmid restriction capabilities of CcSir2/Ago system in E. coli strain

BL20-AI: top -
892
comparison of cell viability in the presence or absence of plasmid-borne CcSir2/Ago expression. Bottom - comparison
893
of plasmid transformation efficiencies in the presence or absence of plasmid-borne CcSir2/Ago expression. C,
894
Quantification of transformation efficiencies for pCDF plasmid with CcSir2/Ago system (three independent replicates,
895
the red line represents average transformation efficiency). D, Cell viability control of BL21-AI, expressing GsSir2/Ago
896
and mutants. E, Control for Fig. 2E - cells contain the pCDF plasmid, however, expression of the GsSir2/Ago system is
897
not induced. F, Qualitative evaluation of pCDF plasmid transformation efficiency in E. coli cells carrying GsSir2/Ago
898
mutants (GsSir2APAZ/Ago, GsSir2/AgoMID and GsSir2/AgoPIWI) of the putative surface of the interaction with nucleic
899
acids. G, Expression analysis of GsSir2/Ago assayed by Western blot. Top: semiquantitative Western blot of the wt
900
GsSir2/Ago complex and its mutants. Numbers above the lanes indicate which part of control protein amount is loaded.
901
The red star shows the lane where the His-tag is on the C-terminus of Ago, rather than the N-terminus of Sir2. GAPDH,
902
loading control. Bottom: Expression analysis of GsSir2/Ago mutants of the putative surface of the interaction with
903
nucleic acids. Three replicates.
904 905
.
CC-BY-NC-ND 4.0 International license
made available under a
(which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is The copyright holder for this preprint
this version posted August 25, 2022.
;
https://doi.org/10.1101/2021.12.14.472599
doi:
bioRxiv preprint
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has

granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
30
A
C
B
D
E
F
G
H
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
Extended Data Fig. 3. Purification and characterization of Sir2/Ago complexes. A, MS analysis of wt GsSir2/Ago complex. The theoretical Mw of the GsSir2 and GsAgo proteins (without 1st Met) are 67929.51 Da and 53135.61 Da, respectively. B, CD spectra of wt GsSir2/Ago and mutants. Mutant spectra are similar to that of a natively folded protein. C, Size-exclusion chromatography of wt GsSir2/Ago, showing elution volume and comparing to mass standards. According to mass spectrometry of the purified GsSir2/Ago complex, the molar mass of the GsSir2/Ago heterodimer is 121 kDa. D, SDS-PAGE analysis of fractions containing the CsSir2/Ago complex eluted from Heparin column. Densitometric inspection shows that Sir2 and Ago proteins are in the ratio ~1:1. Single replicate. E, SDS-PAGE analysis of the CsSir2/Ago stock after dialysis against a storage buffer. Various amounts of the stock solution were loaded on the gel. Densitometric inspection shows that Sir2 and Ago proteins are in the ratio ~0.3:1. Single replicate. F, MS analysis of the CsSir2/Ago complex. The experimental masses (53174.71 Da and 67629.21 Da) are close to the theoretical molecular masses of the Ago protein (53173.91 Da) and the Sir2 protein with the truncated tag at the N terminus (67626.24 Da). G, SEC-MALS analysis of the CcSir2/Ago complex. The experimental mass of 124 kDa is close to the theoretical molecular mass of the CcSir2/Ago heterodimer (121 kDa). H, MS/MS calibration curve of NAD + standard (marked in black) and the observed amount of NAD + (marked in red) in the CcSir2/Ago complex (20 pmol according to the Ago protein). The discrepancy between the expected amount of NAD + (20 pmol, marked in
green) and the actual amount (6.45 pmol, marked in red) was due to the decrease of the Sir2 protein in the CcSir2/Ago
922
preparation (see E).
923 924

.
CC-BY-NC-ND 4.0 International license
made available under a
(which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is The copyright holder for this preprint
this version posted August 25, 2022.
;
https://doi.org/10.1101/2021.12.14.472599
doi:
bioRxiv preprint
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
32
Extended Data Fig. 4. SAXS data. A, Scattering data on an absolute scale. Linear Guinier plot of the initial part of the scattering curve is in the insert. Points cut from the further processing are shown with empty black symbols. B, Kratky plot, normalized by Rg and I(0) parameters. C, Pair distance distribution function. D, CRYSOL Fit of the scattering curve calculated from the GsSir2/Ago AlphaFold model (black curve) with SAXS data (red points).
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
33
A
RsAgo (5AWH)
TtAgo (4N41)
MpAgo (5UX0)
B
CcSir2 dom.
PgSir2 dom.
BcThsA Sir dom.
931
Extended Data Fig. 5. Structural analysis. A, Comparison of GsSir2/Ago, CcSir2/Ago and PgSir2-Ago AlphaFold models with the X-ray structures of long pAgos. Structures are coloured by domains, schematic domain architecture is given above each structure. Guide RNA and target DNA strands are coloured red and blue, respectively. PDB ID codes for long pAgo structures are given in parentheses. RsAgo represents long-B group, TtAgo - longA group of long pAgos (based on Ryazansky et al. classification 8). MpAgo has a distinct OH-type MID domain that is specific to 5'-OH instead of phosphate (MID domain classification - Ryazansky et al.8, MpAgo MID domain biochemical assay - Kaya et al.5). In Sir2/Ago models the N, L1 and L2-like domains previously identified as the APAZ domain correspond to the analogous domains of long pAgos. B, Gs, Cc and Pg Sir2 structural models (cut from full-length models) compared to canonical Sir2 deacetylate TmSir2 and the Sir2 domain of Theoris defence system protein ThsA. Structures are coloured based on secondary structure. Positions corresponding to ThsA Sir2 N112 and H152 are indicated in green. These residues have been shown to be critical for NAD + hydrolysis in ThsA 24. GsSir2 D230 and corresponding positions in other structures are indicated in cyan. NAD + was also superimposed on the ThsA Sir2 structure from TmSir2.
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The

copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
34
A
B
C
D
E
F
Extended Data Fig. 6. EMSA and nucleic acid cleavage experiments. A-B, Binding of single- and double-stranded oligonucleotides by wt GsSir2/Ago and wt CcSir2/Ago, respectively. A radiolabelled strand indicated by the asterisk. C, Binding of complementary DNA targets by GsSir2/Ago binary complexes pre-loaded with RNA guide containing 5'-phosphate terminus in the presence or absence of heparin. To show that no displacement of the radiolabelled guide by the target strand is observed, a control (Cg**) equivalent to the experimental lane marked by a black triangle, but with the guide, rather than the target, bearing the radioactive label, was performed. D, Control EMSA experiments of ssRNA guide binding by wt GsSir2/Ago (left) and non-complementary DNA target binding by the binary wt GsSir2/Ago-gRNA complex in the presence and absence of heparin. E, Representative binding fit curves of several independent replicates used to calculate K_d of ssRNA guide binding by wt GsSir2/Ago (left) and of target DNA binding by wt GsSir2/Ago-gRNA complex (right). F, (No) cleavage activity of various DNA and RNA oligonucleotides by wt GsSir2/Ago and wt CcSir2/Ago. Reaction products were resolved on a 21% denaturing polyacrylamide gel. In heteroduplexes, the asterisk indicates the radiolabelled strand. For panels A-D and F, at least three independent replicates were performed for each experiment.
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
35
961
A
B
C
D
Extended Data Fig. 7. The GsSir2/Ago complex binds NAD + and causes its depletion. A, Two ion transitions were used to detect NAD + in the analysed samples: 662.1 → 540.1 and 662.1 → 426.0. B, MS/MS calibration curve of NAD + standard (marked in black, two replicates) and the observed amount of NAD + in two samples: 93.7 pmol in the wt GsSir2/Ago sample (marked in green), 0.7 pmol in the sample D230A (marked in red). Black dots represent the means of two replicates and error bars are the standard deviation. C, Detection of NAD +. Comparison of the extracted LC-MS/MS chromatograms: ion transition 662.1 → 426.0 of wt GsSir2/Ago sample (panel 1) and NAD + standard (panel 3); ion transition 662.1 → 540.1 of wt GsSir2/Ago sample (panel 2) and NAD + standard (panel 4). Green curves - wt GsSir2/Ago sample, grey curves - NAD + standard. D, Mass chromatogram. For NAD + detection, an extracted ion current (EIC) for [M-H]- m/z = 662.1018 was used. The comparison of EIC signals shows that the amount of NAD + in the samples of the non-induced GsSir2/Ago system in the absence (panel 1) and presence (panel 2) of pCDF plasmid is almost the same, while a significant decrease is observed in the sample of the induced GsSir2/Ago system in the absence of pCDF plasmid (panel 3) and only traces of NAD + are detected in the presence of pCDF plasmid (panel 4).
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The

copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Short prokaryotic Argonautes provide defence against incoming mobile genetic elements through NAD depletion
Mindaugas Zaremba1*, Donata Dakineviciene1, Edvardas Golovinas1, Evelina Zagorskaite1, Edvinas Stankunas1, Anna Lopatina2, Rotem Sorek2, Elena Manakova1, Audrone Ruksenaite1, Arunas Silanskas1, Simonas Asmontas1, Algirdas Grybauskas1, Ugne Tylenyte1, Edvinas Jurgelaitis1, Rokas Grigaitis1, Kstutis Timinskas1, eslovas Venclovas1, Virginijus Siksnys1*

Supplementary Information

Contains Supplementary Note, Supplementary Methods, Supplementary References, and 3 Supplementary Tables
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

987 Supplementary Note 1
Bioinformatic analysis of the selected Sir2/Ago systems. Closer inspection of the genomic neighbourhood of the selected pAgos showed that no other conserved operons are formed with the pAgo genes. However, the observed enrichment of putative restriction endonucleases, mobile genetic elements (e.g. transposases, integrases) and toxin-antitoxin systems in the neighbourhood of the GsAgo and CcAgo genes indicates that they could be a part of so-called bacterial defence islands 1. All three systems have MID-PIWI domains, characteristic of Agos, but their PIWI domains are catalytically inactive (Fig. 1A). Importantly, this derived feature is shared with the evolutionary closest group of long pAgos (Extended Data Fig. 1C). Although overall sequence similarity between the MID-PIWI regions of selected systems (Gs, Cc and Pg) and long pAgos is rather low (~15-20% sequence identity), their MID domains do have conserved residues associated with binding of the 5'-end of the guide strand 2 (Extended Data Fig. 1E). The conserved motif of MID domains of all three systems is of the HK-type, which is more similar to the 5'-P-end binding motif present in long pAgos such as RsAgo or TtAgo than to the 5'-OH-end binding motif found, for example, in MpAgo 2. Unlike long pAgos, short pAgos lack the PAZ domain and are rather associated with an APAZ (analogue of PAZ) domain (Fig. 1A)3. By phylogeny of the APAZ domain region, PgSir2-Ago belongs to Ia, and GsSir2/Ago and CcSir2/Ago to Ib groups, respectively (Extended Data Fig. 1B)2. The sequences of Sir2 domains associated with the three studied pAgos are similar to canonical sirtuins 3, and also have an identifiable signature of a putative NAD +-binding pocket (Extended Data Fig. 1C). Yet there are many differences from canonical sirtuins, and the identified conserved amino acid positions found only in pAgo-associated Sir2 domains suggest that these domains might have a function different from the typical deacetylase activity of sirtuins (Extended Data Fig. 1C). pAgos-associated Sir2 domains also show similarities to NADase domains of ThsA proteins from the Thoeris anti-phage systems (Extended Data Fig. 1C)4,5.
GsSir2/Ago complex resembles long PAZ-free pAgos containing an additional effector domain. To get an insight into the 3D structure of GsSir2/Ago and the other two systems, we used AlphaFold 6 to generate corresponding structural models. Based on high AlphaFold confidence values (plDDT>87; pTM>0.82) and favourable VoroMQA 7 statistical energy scores (>0.53) the models for all three systems might be expected

to be of comparable quality with experimental structures. Consistent with clear homology between the three systems the corresponding models showed close structural similarity with each other (Extended Data Fig. 5). Therefore, we focus only on the GsSir2/Ago complex. In this complex, the Sir2 and Ago subunits bind together to form a structure similar to that of a single-chain long pAgo protein such as RsAgo (Extended Data Fig. 5). Based just on a structural similarity search against PDB, GsSir2/Ago appears to be most similar to the TtAgo structure classified as a long-A pAgo 2. Another close structural match is RsAgo, a member of long-B pAgos 2. GsAgo shares 17% identical residues with both TtAgo and RsAgo. However, since RsAgo, just like short pAgo proteins, has a catalytically inactive PIWI domain, a comparison of GsSir2/Ago with RsAgo might be biologically more relevant.

The APAZ part of the GsSir2 chain is structurally similar to the combination of N, L1 and L2 domains of RsAgo (similarity to the N domain has been already proposed earlier 8,9). However, the L2 linker domain in GsSir2/Ago corresponds to the two fragments, C-terminus of Sir2 and N-terminus of Ago. Importantly, the PAZ domain, required for the 3'-end recognition of the guide strand in long pAgos, is missing from the GsSir2/Ago structure altogether. Additionally, GsSir2/Ago has a smaller N domain than long pAgos, which could also alter nucleic acid binding. PIWI and MID domains are structurally similar to corresponding RsAgo domains. Notably, GsSir2/Ago has a longer loop (residues 268-272) compared to the corresponding region in RsAgo.

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

This loop shows some steric overlap with the copied-in RNA/DNA duplex, but presumably, its conformation may adjust to accommodate RNA/DNA heteroduplex. The loop contains two positively charged residues (R269, K270) that might be involved in the binding of the nucleic acid backbone.

In the structural model, the N-terminal Sir2 domain of the GsSir2 protein is attached to the C-terminal domain corresponding to the N-terminal region of a long pAgo through a long linker. In effect, the two domains of GsSir2 are positioned at the opposite extremes of the complex (Sir2 domain is bound to the PIWI domain whereas the C-terminal domain is bound to the MID domain). Sir2 domain of GsSir2/Ago is structurally similar to canonical Sir2 proteins suggesting it could bind NAD + in a similar way. However, the overall structurally closest homolog is the Sir2 domain of the ThsA protein from the Thoeris defence system 5 (Extended Data Fig. 5). The ThsA residues, N112 and H152, shown to be essential for NAD + hydrolysis have their counterparts in GsSir2 (N142 and H186). Interestingly, the loop containing H186 in different GsSir2/Ago models displays some conformational heterogeneity hinting at possible flexibility, which might be relevant for the activity regulation. Conservation of these two positions is also observed in other Sir2 homologs (Extended Data Fig. 1C). On the other hand, the GsSir2 D230 residue shown here to be important for NAD + hydrolysis is conserved in ThsA Sir2 domain, but not in the canonical Sir2 proteins (e.g., V193 in TmSir2). In the AF2 structural model, the NAD +-bound Sir2 active site is hidden at the interface between the MID and PIWI domains. One may assume that binding of the complementary DNA target by the binary GsSir2/Ago-gRNA complex triggers a conformational change that opens and activates the Sir2 active site resulting in NAD + hydrolysis.

The activated GsSir2/Ago depletes NAD + enzymatically. Our results show that wt GsSir2/Ago tightly interacts with NAD + (Fig. 5A, Extended Data Fig. 7). Therefore, there is a theoretical possibility that the NAD + depletion in E. coli cells expressing the GsSir2/Ago system might be related to its tight binding by the activated GsSir2/Ago complex. However, this assumption can be ruled out based on our approximate calculations of relative levels of the GsSir2/Ago protein and NAD + molecules in E. coli cells. According to our semiquantitative Western Blot analysis (Extended Data Fig. 2G), the GsSir2/Ago concentration (~1 M) is more than ~1000-fold lower than that of NAD + (~2.5 mM), which is in the range of the published NAD + concentrations (~0.6-8 mM)10. Therefore, the activated GsSir2/Ago complex should

act enzymatically to deplete such a large excess of NAD +.
1065 Supplementary methods
Sequences analysis. Lists of pAgo homologues and associated APAZ domains were retrieved from supplementary data of the Ryazansky et al. article 2. Thermotoga maritima Sir2 homologues were collected from the SwissProt database 11 using BLAST 12 (1e-5 e-value cutoff). Full-length sequences were retrieved from NCBI. Bacillus cereus ThsA (PDB ID 6LHX) homologues were collected using BLAST from UniRef 50}^{13 database (1e-10 e-value cutoff). To remove other Sir2 homologues, these sequences were clustered with CLANS 14. The cluster separated at p=1e-70 was selected as the representative ThsA group. Fragmented ThsA sequences and sequences missing one of the domains were discarded. Multiple sequence alignments were generated using MAFFT (l-INS-i mode for high accuracy)15. Jalview 16 was used for multiple sequence alignment analysis, cutting and visualization. Sequence motif visualization was done using WebLogo 3 server 17. Construction of combined multiple sequence alignment of GsSir2, TmSir2 and ThsA homologues was guided by the alignments between GsSir2, TmSir2 and BsThsA sequences obtained using the HHpred server 18.
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
39
Phylogenetic tree construction. The phylogenetic trees were constructed with FastTree 19 using the WAG model of amino acid substitution 20 and the gamma model of rate heterogeneity. Prior to phylogenetic analysis, multiple sequence alignment positions containing more than 50% gaps were removed using trimAl 21. Phylogenetic tree visualization was done with iTOL 22.
Genomic neighbourhood analysis. For genomic neighbourhood analysis, the Geobacter sulfurreducens, Caballeronia cordobensis, and Paraburkholderia graminis genomes (respective GenBank accessions: GCA_000210155.1, GCA_001544575.2 and GCA_000172415.1) and all associated sequence and annotation data were obtained from NCBI (ftp://ftp.ncbi.nlm.nih.gov/genomes/Bacteria/). Genes in the neighbourhood of each pAgo (10 upstream and 10 downstream) were identified based on available genome annotations. To refine available functional annotations of these genes, searches through Pfam 23 and PDB databases were performed using the HHpred server 18.
1091 Expression and purification of CcSir2/Ago complex. Expression vector constructs of the 1092 CcSir2/Ago system were used to transform E. coli BL21(DE3) strain. Transformed bacteria were 1093 grown at 37 ° C in LB medium in the presence of 50 g/ml ampicillin until OD600 = 0.7 was reached. Then, the medium was cooled to 16 ° C temperature and proteins were expressed for 16 h by adding 0.1% w/v L-arabinose. Harvested cells were disrupted by sonication in buffer A (20 mM Tris-HCl (pH 8.0 at 25 ° C), 1.0 M NaCl, 2 mM phenylmethylsulfonyl fluoride, 5 mM 2-mercaptoethanol), and cell debris was removed by centrifugation. TwinStrep-CcSir2/Ago complex was purified to > 90% homogeneity by chromatography through Strep-Tactin XT Superflow (Iba), HiLoad Superdex 200, HiTrap Heparin HP columns (GE Healthcare). Purified proteins were stored at -20 ° C in a buffer containing 20 mM Tris-HCl (pH 8.0 at 25 ° C), 500 mM NaCl, 2 mM DTT and 50% v/v glycerol. The identity of the purified proteins was confirmed by mass spectrometry. Protein concentrations were determined from OD280 measurements using the theoretical extinction coefficients calculated with the ProtParam tool available at http://web.expasy.org/protparam/. TwinStrep-CcSir2/Ago complex concentrations are expressed in terms of heterodimer.
CD analysis. To test whether the introduced mutations have an effect on secondary structures of the GsSir2/Ago, CD spectra of the proteins were recorded. Protein samples (5 M of heterodimer) were prepared in 200 l of the buffer (10 mM Tris-HCl (pH 8.0 at 25 ° C), 50 mM NaCl). The circular dichroism spectra were recorded in triplicate with Jasco J-815 CD spectrometer (Jasco, Easton, MD) using a 1 mm quartz cuvette at a scanning speed of 50 nm/min from 200 to 250 nm. The temperature of samples was controlled at 25 ° C using a Jasco temperature control device.
Construction and analysis of Sir2/Ago structural models. Structural models were generated using the

AlphaFold method 6 implemented as ColabFold 24, an online Google Colaboratory notebook. For modelling, the ‘Alphafold2_advanced’ notebook (colab.research.google.com/github/sokrypton/ColabFold/blob/main/beta/AlphaFold2_advanced.ipy nb) was used. GsSir2/Ago and CcSir2/Ago complexes were modelled as heterodimers, whereas the Pg sequence representing a fusion of Sir2 and a short pAgo (PgSir2-Ago) was modelled as a monomer. The modelling pipeline was run with default parameters except for the multiple sequence alignment (MSA) pairing, which was set to ‘paired+unpaired’. The best-of-five model in each case was selected using VoroMQA 7. Structure similarity searches of models/domains against PDB were performed using Dali 25. Structure analysis and visualization was performed using UCSF Chimera 26. Putative binding sites of Sir2/Ago complexes were investigated by simply copying the RNA/DNA duplex and NAD + from RsAgo (PDB ID: 5AWH) and TmSir2 (PDB ID: 2H4F) structures correspondingly after their superposition onto structural models. No attempts to remove possible clashes between protein models and either RNA/DNA or NAD + were made.

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Nucleic acid cleavage assay. The same linear oligonucleotides used for EMSA (Supplementary Table 2) were given as substrates for nucleic acid cleavage assay. To raise the total substrate concentration, 5'-32P radiolabelled oligonucleotides were mixed with appropriate cold 5'P-oligonucleotides at a ratio of 1:4 of hot:cold, and diluted to a working concentration of 100 nM in reaction buffer (33 mM Tris-acetate, pH 7.9, supplemented with 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA, 5 mM DTT). Protein dilutions were made using the same reaction buffer to 2x final reaction concentration. Protein complexes and nucleic acids were mixed in final concentrations of 50 nM NAs and 0, 50 or 500 nM protein, incubated for 1 hour at 25 ° C. The reaction was stopped with the addition of 2x 95% formamide dye and incubating for 5 min at 95 ° C. Reaction products were resolved by denaturing PAA gel electrophoresis (21% 29:1 acrylamide/bis-acrylamide in TBE (89 mM Tris, 89 mM boric acid, 2 mM EDTA), supplemented with 8 M urea), visualised with a phosphor imager and analysed in OptiQuant software.

1137

1138

bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

1139 Supplementary tables

1140 Supplementary Table 1. Strains, bacteriophages, plasmids, and proteins used in this work

Bacterial strains (Escherichia coli)	Details	Source or reference, links
MG1655	F^{-} \lambda^{-} ivG^{-} rfb-50 rph-1	Blattner, F. et al.^{28
BL21-Al	F^{-} ompT hsdS_{B}(r_{B}m_{B}) gal dcm araB::T7RNAP-tetA	Bhawsinghka, N. et al.^{29
DH10B	F^{-} mcrA \Delta(mrr-hsdRMS-mercBC) \Phi 80dlacZ\Delta M15 \Delta acX74 \text{ endAl recA1 deoR} \Delta(ara,leu)7697 \text{ araD139 galU galK nupG rpsL} \lambda^{-	Grant, S.G.N. et al.^{30
Bacteriophages (Escherichia coli)		
T7	NC_001604.1	Udi Qimron; Kulezyk, A. & Richardson, C.C. 31
lambda-vir	NC_001416.1	Udi Qimron; Casjens, S.R. & Hendrix, R.W. 32
SECphi27	LT961732.1	Sorek lab collection; Doron, S. et al.^{4
SECphi18	LT960607.1	Sorek lab collection; Doron, S. et al.^{4
T4	AF158101.6	Udi Qimron; Yap, M.L. & Rossmann, M.G. 33
SECphi17	LT960609.1	Sorek lab collection; Doron, S. et al.^{4

Plasmids		
pBAD/HisA	Bacterial expression vector, carrying a pBR322 replicon and ampicillin resistance, provides an N-terminal His_{6 tag for expressed proteins, <40 copies/cell.	ThermoFisher cat#V43001. Tolia, N.H. & Joshua-Tor, L.^{34 https://benchling.com/s/seq-tfRnXVmxwmGJqYnXmsDw?m=slm-gnSb1LLWrQibTPI2ibbs
pBAD24	Bacterial expression vector, carrying pBR322/ColE1 and F1 replicons, and ampicillin resistance gene, ~40 copies/cell.	Guzman, L.M et al.^{35 https://benchling.com/s/seq-zaoDsxfoymWzOCDoHSEI?m=slm-ALGNMJ5F3aprLKyOrHOK
pBAD24-HSH	pBAD24 vector with an HSH affinity tag before the rrnB terminator. Used to fuse the tag on target protein C-terminus, ~40 copies/cell.	Guzman, L.M et al.^{35
pCDF	Bacterial vector carrying a CloDF13 replicon, lacI gene and streptomycin/spectinomycin resistance gene, 20-40 copies/cell.	Tolia, N.H. & Joshua-Tor, L.^{34 https://benchling.com/s/seq-B4oyRceI17nWdhFpsWmGx?m=slm-SrF4zHJbfBfIhrG49jP
pCDF_Kn	pCDF vector, carrying kanamycin resistance instead of the canonical streptomycin, 20-40 copies/cell.	In-house construct.34 https://benchling.com/s/seq-7KKOFJRtBzzCbjA4MN3L?m=slm-Ez8cisdT8LtXDMSNSIur
pCOLA	Bacterial vector with a ColA origin of replication, harbours kanamycin resistance gene, 20-40 copies/cell.	Tolia, N.H. & Joshua-Tor, L.^{34 https://benchling.com/s/seq-dtqZ232iXhuGqGiyZR0a?m=slm-hLBJqu72mbC0yEkkFKeu
pACYC184	Bacterial vector carrying p15A origin of replication and tetracycline resistance gene, 10-12 copies/cell.	Tolia, N.H. & Joshua-Tor, L.^{34 https://benchling.com/s/seq-Rz0xd8JhIeZa1vwsFx7c?m=slm-5rwfx80s5z1sp04YeEbJ
pRSF	Bacterial vector carrying the RSF1030 replicon, lacI gene and kanamycin resistance gene, >100 copies/cell.	Tolia, N.H. & Joshua-Tor, L.^{34 https://benchling.com/s/seq-7C0lDtWUXncuAQyKn19F?m=slm-wTVKOxRQQYAFlv1CXJET
pCDF_ColA	pCDF plasmid with its ori site replaced with the ori site from a pCOLA plasmid.	In-house construct.34 https://benchling.com/s/seq-yXHSYIuEhly5tKTDgHmb?m=slm-PQajuJX3EDmQqYnXOw9k
pCOLA_CloDF13	pCOLA plasmid with its ori site replaced with the ori site from a pCDF plasmid.	In-house construct.34 https://benchling.com/s/seq-PGM8cj3vsbJ2foudD4gS?m=slm-CUFdjxao5BVO2rDYJ9w
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

pBAD/HisA_GsSir2/Ago	Bacterial expression vector with Sir2 and Ago genes from G. sulfurreducens.	https://benchling.com/s-seq-6CDMPX5NCY2UZSExH9YR?m=slm-sDqIpXnJdAHVagPVIDcR
pBAD/HisA_GsSir2(D230A)/Ago	Bacterial expression vector with a D230A mutational variant of Sir2 and Ago genes from G. sulfurreducens.	https://benchling.com/s-seq-8xPH4KTJbHyGjfhzx2KS?m=slm-SNhTzzHe0Wen4MPLBNoA
pBAD24_GsSir2/Ago-HSH	Bacterial expression vector with Sir2 and Ago genes from G. sulfurreducens, with an HSH affinity tag on Ago protein C-terminus.	https://benchling.com/s-seq-kzi2cWwSY2P2Gc2wV6AFAF?m=slm-CFvHs1oLhUQY5NcM3MyU
pBAD/HisA_TwinStr-ep_TEV_GsSir2/Ago	Bacterial expression vector with an N-terminal His6-TwinStrep-TEV-carrying Sir2 and an Ago protein gene from G. sulfurreducens.	https://benchling.com/s-seq-1tCIsHyBFY7ba7Ummul?m=slm-v7BRSSk20CZ8e94uXn9p
pBAD/HisA_TEV_CcSir2/Ago	Bacterial expression vector with an N-terminal His6-TEV-carrying Sir2 and an Ago protein gene from C. cordobensis.	https://benchling.com/s-seq-3YCAQfAp34mRvXctlpkb?m=slm-6GkVFazYraa8IXQCYF6r
pBAD/HisA_TwinStrep_TEV_CcSir2/Ago	Bacterial expression vector with an N-terminal His6-TwinStrep-TEV-carrying Sir2 and an Ago protein gene from C. cordobensis.	https://benchling.com/s-seq-uHOLRoSabse4I5PN5jR5?m=slm-EcbYQvycbLNVXc3uEVte

|pBAD24_PgSir2-Ago|Bacterial expression vector with a Sir2-Ago fused protein gene from P. graminis.|https://benchling.com/s/seq-oLr3DRXdEfJR2W7tWNdP?m=slm-qAwOAfwDnuHTQaaINYEQ|
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

1142 Supplementary Table 2. Oligonucleotides used in this work.
Oligonucleotide	Sequence (5'->3' direction)	Description
Oligonucleotides used for cloning, mutations and controls		
MZ-239	GCTGATGGTCTCGCTAGCATGGATGTCTTAACTGACAATGAGTTTTAC	Forward primer for the amplification of the GsSir2/Ago operon.
MZ-240	CTGATTCTCGAGTTACATAAAAACCGATAATCATATATTCGTTAAC	Reverse primer for amplification of the GsSir2/Ago operon.
MZ-351	GTGGTCGTGCTGAAAATGTTATGACTATGCTTC	Mutagenic primer (top strand) for the D230A mutation within the GsSir2 protein.
MZ-352	GTCATAACATTTTCAGCACGACCACTATAGCC	Mutagenic primer (bottom strand) for the D230A mutation within the GsSir2 protein.
MZ-325	GCTGATGGTCTCGCATGGATGTCTTAACTGACAATGAGTTTTAC	Forward primer for the amplification of the GsSir2/Ago operon.
MZ-326	TGATGCTCGAGCATAAAAACCGATAATCATATATTCGTTAAC	Reverse primer for the amplification of the GsSir2/Ago operon.
MZ-915	TGATTACCTGCTTAGCATGCATAATCGCGCTGCTTGCAAACG	Forward primer for the amplification of the PgSir2-Ago gene.
MZ-916	TGATTAAGCTTTACATAAACAGACGGTAATCGTACGAGTGGTC	Reverse primer for the amplification of the PgSir2-Ago gene.
MZ-1217	GAAATCTTAGAGTGATGGTGTCGGGAATCC	Forward primer for the amplification of the COLA ori.
MZ-1218	AGTGATGGTGTCGGGAATCC	Reverse primer for the amplification of the COLA ori.
MZ-1230	AAATAGCTAGCTCACTCGGTC	Forward primer for the amplification of the CDF ori.
MZ-1231	GAAATCTAGAGCGGTTCAGTAG	Reverse primer for the amplification of the CDF ori.
TS-48	CAUGCUGCAGGUCGACUUG	22 nt RNA oligonucleotide used as a control analysing GsSir2/Ago-associated nucleic acids.
Oligonucleotides used for EMSA and cleavage experiments		
ssRNA (TF-A)	AUAAUGGUUUCUUAGACGUCGUUUUAGAGCUGUGUUUGUUCG	42 nt RNA oligonucleotide
ssDNA (TK-49)	ATAATGGTTTCTTAGACGTCGTTTTAGAGCTGTGTTGTTCG	42 nt DNA oligonucleotide, sequence identical to TF-A, only DNA.
ssDNA (MZ-949)	CGAAACAACACAGCTCTAAAACGACGTCTAAGAAACCATTAT	42 nt DNA oligonucleotide, complementary to TF-A and TK-49
nsp-ssDNA (MZ-589)	GGAGATGGGAGATCATTAGATGTAGAACTTGAATGGCAACGG	42 nt DNA oligonucleotide, non-complementary to TF-A
ssRNA (MZ-1185)	CGAAACAACACAGCUCUAAAACGACGUCUAAGAAACAUAU	42 nt RNA oligonucleotide, complementary to TF-A and TK-49
dsRNA	5' -AUAAUGGUUUCUUUGACGUCGUUUUAGAGCUGUGUUUUGC-3';3' -UAUACCAAAGAUCUGCAGCAAAAUCUGACACAAACAAAGC-5'	TF-A annealed to MZ-1185
dsDNA	5' -ATAATGGTTTCTTAGACGTCGTTTTAGAGCTGTGTTGTTTTCG-3';3' -TATTACAAAGAATCTGCAGCAAAATCTCGACACAAACAAAGC-5'	TK-49 annealed to MZ-949
RNA/DNA	5' -AUAAUGGUUUCUUUGACGUCGUUUUAGAGCUGUGUUUUGC-3';3' -TATTACAAAGAAATCTGCAGCAAAATCTCGACACAAACAAAGC-5'	TF-A annealed to MZ-949
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The

copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
44
1145 Supplementary Table 3. Parameters of the SAXS data.
Recipient space parameters	Merged SAXS normalized against water scattering merged data
s \min (nm^{-1}) as in Guinier range	0.1116200
Guinier range (points) used for Guinier analysis	48 to 170
Rg, by Guinier analysis / AUTORG, nm	3.61 / 3.61 \pm 0.034
I(0) AUTORG, (cm^{-1})	0.0734 \pm 0.00022
Rg*s by Guinier analysis	1.33281
Molecular Mass estimations out of SAXS data, kDa	
PRIMUS Qp	107.9
PRIMUS Vc	99.7
PRIMUS Size&Shape	125.3
PRIMUS Absolute scale	101.1
PRIMUS Bayesian inference	101.1
SAXSMoW server, v2.1	124.8
Real-space parameters (GNOM)	
s range (nm^{-1}) used in GNOM	0.1116 to 3.2449
Dmax as parameter in GNOM / DATCLASS / SHANUM, nm	11.4 / 12.3 / 13.4
Rg real GNOM / DATPOROD, nm	3.579 \pm 0.005044 / 3.58
I(0) real GNOM / DATPOROD, (cm^{-1})	0.07277 \pm 0.0001016 / 0.0728
Porod volume DATPOROD (nm^{3})	185.06
SASDB ID	SASDNH2
1146
1147
References
References
bioRxiv preprint doi: https://doi.org/10.1101/2021.12.14.472599; this version posted August 25, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.
47
1225 Methods 3, 55–64 (2006).
Guzman, L. M., Belin, D., Carson, M. J. & Beckwith, J. Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. J. Bacteriol. 177, 4121–4130 (1995).
1229
1230
1231